

Provided for non-commercial research and education use.

Not for reproduction, distribution or commercial use.



This article was published in an CASRP journal. The attached copy is furnished to the author for non-commercial research and education use, including for instruction at the authors institution, sharing with colleagues and providing to institution administration.

Other uses, including reproduction and distribution, or selling or licensing copied, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding CASRP's archiving and manuscript policies encouraged to visit:

<http://www.casrp.co.uk/journals>

© 2015 CASRP Publishing Company Ltd. UK.



Available online at www.casrp.co.uk/journals

CASRP PUBLISHER
International Journal of Advanced Biological
and Biomedical Research 3(4) (2015) 378–383

doi: 10.18869/IJABBR.2015.378



Original Article

Open Access

Effects of different levels of nano chromium chloride in diet on egg quality and blood chromium content of layingjapanese quail

M. Amiri Andi*, A. Shahamat

Department of Animal Science, Sanandaj Branch, Islamic Azad University, Sanandaj, Kurdistan, Iran.

Abstract

The goal of this experiment was to investigating of effects of different levels of nano chromium (N-Cr) in diet on egg quality and blood chromium content of japanese quail. Two hundred and forty laying japanese quail (45 day old) were randomly distributed to 20 experimental unit with 12 birds in cage pens of 40 × 80 cm. Experimental treatments include: 1) control national research council (NRC, 1994) based diet (with no N-Cr), 2) 200 ppb, 3) 400 ppb, 4) 600 ppb and 800 ppb of N-Cr. Egg weight, egg internal quality (Huagh unit) and blood chromium were

*Corresponding author: Department of Animal Science, Sanandaj Branch, Islamic Azad University, Sanandaj, Kurdistan, Iran.

© 2015 The Authors. This is an open access article under the terms of the Creative Commons Attribution-Non Commercial- No Derives License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Received 12 November 2015

Accepted 13 December 2015

Available online 19 December 2015

iThenticate screening 16 November 2015

English editing 11 December 2015

Quality control 15 December 2015

determined. Treatments influenced egg weight, yolk weight, albumen height, albumen weight, shell weight and thickness and Haugh unit ($p \leq 0.05$). But blood chromium content was not affected by levels of N-Cr ($p \geq 0.05$). In conclusion, it seems that N-Cr level of 800 ppb had further effects on Japanese quail egg quality.

© 2015 Published by CASRP publishing company Ltd. UK. Selection and/or peer-review under responsibility of Center of Advanced Scientific Research and Publications Ltd. UK.

Keywords: Japanese quail, Nano chromium, Egg quality, Haugh unit, blood chromium.

1. Introduction

Chromium (Cr) is an essential mineral element for livestock (Lukaski, 1999). Notwithstanding, national research council (NRC, 1994) had no recommendation for Cr requirement of poultry especially for Japanese quail. But, in recent years, more researches in using of Cr and its useful effects on poultry egg and meat quality were done (Sahin et al., 2005; Sahin et al., 2002; Lien et al., 1999). Supplemental Cr in the form of Cr-picolinate (1.2 mg/kg) resulted in further egg weight of quail (Sahin et al., 2001). Derya et al., (2009) demonstrated that supplemental Cr-picolinate had increasing effect on shell thickness but no effect on egg weight of Japanese quail. In addition, with adding supplemental Cr to diet, egg weight, shell thickness and Haugh unit were linearly increased (Sahin et al., 2002). Today, nanotechnology prepared new Cr resources named N-Cr that have further bioavailability than organic and inorganic Cr supplements (Zha Ly et al., 2008). Positive effects of N-Cr on performance, egg quality and tissues mineral accumulation in layer chickens were shown by Sirirat et al., (2013). They use three levels of nanoparticles Cr-picolinate (0, 500 and 3000 ppb) in their experiment. Chromium chloride (CrCl_3) used in poultry nutrition (Eren and Baspnar, 2004) but not in nano form. Nano chromium (N-Cr) in chemical form of chromium chloride were not used in laying Japanese quail diet, yet. Therefore, the goal of this experiment were to investigate effects of different levels of nano chromium (N-Cr) in diet on egg quality and blood chromium content of laying Japanese quail.

2. Materials and methods

Two hundred and forty laying Japanese quail (45 day old) were randomly distributed to 20 experimental units with 12 birds in cage experimental units of 40×80 cm, in a completely randomized design for 7 weeks. The five treatments used in the present experiment included of the following: (1) basal diet was formulated based on national research council (NRC, 1994) (Table 1); (2) basal diet plus 200 ppb Cr as nanoparticles chromium chloride (N-Cr); (3) basal diet plus 400 ppb Cr as N-Cr; (4) basal diet plus 600 ppb Cr as N-Cr and (5) basal diet plus 800 ppb Cr as N-Cr. The nano chromium chloride was prepared by Industrial Research Institute of Iran (Tehran, Iran). The mean particle size of N-Cr chloride was 80.73 ± 2.96 nm. Feed and water were *ad libitum*. Lighting program was about 16 hours. The quails were reared according to commercial quail production conditions.

2.1. Egg parameters measurement

Total egg produced were daily numbered and weighed with an electronic scale with 0.01 g accuracy. At the end of the experiment four egg per experimental unit were randomly selected for egg traits measurements. Yolk weight, albumen weight and shell weight of each egg was measured with an electronic scale with 0.001 g accuracy. Thickness of egg shell in the equator, blunt end, and pointed end of the eggs (excluding the membrane of shell) was measured with a micrometer (OSK-13469, OGAWA SEIKI CO., LTD. JAPAN). For measuring of Haugh unit, after weighing of each egg, the egg was broken and albumen height were measured by an albumen height determiner (OSK-13471, OGAWA SEIKI CO., LTD. JAPAN). The Haugh unit (HU) was measured:
$$HU = 100 \log (H + 7.57 - 1.7^{0.37})$$

2.2. Blood chromium

On day 93 of trial, blood samples from the quails were collected and placed into trace-mineral-free tubes with sodium heparin, and then centrifuged for 15 min at $3000 \times g$ at 4°C (SIGMA 4-15 Lab Centrifuge, Germany).

After centrifugation, the fresh plasma was collected and analyzed for chromium. Plasma chromium levels were analyzed by using atomic absorption spectrophotometer (PerkinElmer, AA 600, USA).

2.3. Statistical analysis

Data were subjected to statistical analysis using analysis of variance (ANOVA) appropriate for a completely randomized design. When significant effects were detected by ANOVA, treatment means were compared using Duncan's multiple range tests. All statistical analyses were performed with SAS (SAS, 2001). Differences were considered significant at $P < 0.05$. All of parameters were analyzed as follows: $Y_{ij} = \mu + T_i + e_{ij}$, where Y_{ij} =individual observation, μ =overall mean, T_i =effect of treatment and e_{ij} represents the random error.

Table 1
Composition of basal diet.

Ingredients	% Diet
Corn	50.48
Soybean meal	32.36
Vegetable oil	1.92
Meat meal	7
Ca CO ₃	6.89
DCP	0.19
Na ₂ CO ₃	0.14
Salt	0.17
dl-methionine	0.29
Lysine hydro chloride	0.06
Premixe(vitamin ¹ +mineral ²)	0.5
Total	100
Calculated nutrients	
Metabolizable energy (kcal/kg diet)	3000
Crud protein	20
Ca	2.5
P	0.35
Methionine	0.45
Lysine	0.85

¹Supplied per kg of vitamin mixture: Vitamin A: 7.2gr; Vitamin D: 7. gr; Vitamin E: 14.4gr; Vitamin K3: 1.6gr; Vitamin B1: 0.72gr; Riboflavin: 3.3gr, Pantothenic acid: 12gr, niacin: 12160 mg; Vitamin B6: 6.2 mg; Biotin: 0.2 gr; Vitamin B12 - 0.6 gr; choline chloride 440. ² Supplied per kg of mineral mixture: manganese (oxide): 64 gr; iron (FeSO₄) -100 gr; zinc (oxide): 44 gr; copper (CuSO₄):16 gr; iodine (calcium iodate): 64 gr; selenium (1%): 8 gr; cobalt :0.2 gr.

3. Results

3.1. Egg traits

Mean egg traits of laying japanese quail were shown in Table 2. Over all egg traits were influenced by the N-Cr levels of diet. Quails receiving 800 ppb of N-Cr had the greatest significant egg weight than 0, 200 and 400 ppb receiving birds (13.52 vs. 10.60, 11.20 and 11.30 g, respectively) ($p \leq 0.05$). Treatment 5 had the highest yolk weight than treatments 1 and 3 (4.31 vs. 3.73 and 3.80 g, respectively) ($p \leq 0.05$). Albumen height in 800 ppb group was significantly higher than control and 200 ppb group (2.54 vs. 1.58 and 1.66 mm, respectively) ($p \leq 0.05$). Quails receiving 800 ppb of N-Cr had the greatest significant albumen weight than control (7.09 vs. 6.24 g, respectively) ($p \leq 0.05$). Quails in treatment 5 (800 ppb N-Cr) had significant greatest shell thickness and HU than control (387 vs. 310 μ m, for shell thickness and 75.92 vs. 68.59, for HU, respectively).

3.2. Blood chromium

Treatments had no effects on blood chromium. Notwithstanding, quails in 800 ppb group had the highest blood chromium.

Table 2
Effects of different levels of nano chromium in diet on egg quality.

Egg traits	Levels of nano chromium in diet (ppb)					Significance
	0	200	400	600	800	
Egg weight (g)	10.60 ^b ±0.19	11.20 ^b ±0.32	11.30 ^b ±0.41	11.74 ^{ab} ±0.37	13.52 ^a ±1.20	**
Yolk weight (g)	3.73 ^b ±0.18	3.90 ^{ab} ±0.13	3.80 ^b ±0.15	4.10 ^{ab} ±0.09	4.31 ^a ±0.10	*
Albumen height (mm)	1.58 ^b ±0.08	1.66 ^b ±0.09	2.01 ^{ab} ±0.30	2.05 ^{ab} ±0.21	2.54 ^a ±0.18	**
Albumen weight (g)	6.24 ^b ±0.32	6.74 ^{ab} ±0.22	6.63 ^{ab} ±0.25	6.74 ^{ab} ±0.26	7.09 ^a ±0.16	*
Shell weight (g)	1.28 ^b ±0.03	1.42 ^{ab} ±0.04	1.37 ^b ±0.07	1.44 ^a ±0.08	1.56 ^a ±0.02	*
Shell thickness (µm)	310 ^b ±20	344 ^{ab} ±14	341 ^{ab} ±19	355 ^{ab} ±24	387 ^a ±22	*
Huagh unit (HU)	68.59 ^b ±0.47	71.50 ^{ab} ±1.14	71.59 ^{ab} ±2.26	71.21 ^{ab} ±1.54	75.92 ^a ±1.53	*

Data are mean ± SD. a-b: In each raw means with different superscript had significant differences (p≤0.05).

Table 3
Effects of different levels of nano chromium in diet on plasma chromium.

Trait	Levels of nano chromium (ppb)					Significance
	0	200	400	600	800	
Plasma chromium (mg/dl)	221.47±2.54	224.40±2.57	222.15±2.41	222.78±2.10	226.14±3.26	Ns

Ns: not significant. Data are mean ± SD.

4. Discussion

There is no report using nanoparticle chromium in laying quails. Perhaps our experiment is the first one that used nanoparticle chromium chloride. Chromium is a keypoint of glucose tolerance factor (GTF) then it can influencing insulin performance that is a most important anabolic hormones (Anderson et al., 1997; Holdsworth and Neville, 1990). Due to insulin, body cells can use glucose and sufficient amino acids enter to cells, so performance increases (Hossain et al., 1998). Perhaps, with this reason, in our experiment egg weight increased. Our results are parallel to Kim et al., (1997). They used Cr-picolinate in their experiment and concluded 800 ppb of Cr in the diet resulting in further egg weight of laying hens. Level of 800 ppb of N-Cr chloride had highest effect on yolk weight. These conclusions are similar to Sahin et al., (2001) and Yildiz et al., (2004) in laying quails and Uyanick et al., (2002) in laying hens but not to Sirirat et al., (2013). In our experiment the 800 ppb of N-Cr chloride had significant beneficial effects on albumen weight, albumen height and HU. Sahin et al., (2001) and Yildiz et al., (2004) in laying quails and Uyanick et al., (2002) in laying hens were taken same results but Sirirat et al., (2013) was

taken similar result in HU, only. Jensen et al., (1987) reported that Cr had useful effects on the internal quality of egg. It seems, there is a linear effect between supplemental Cr levels in the diet and significant increasing of HU, Sahin et al., (2002) reported. In the experiment of Sirirat et al., (2013), two levels of nano chromium picolinate (500 and 3000 ppb) had beneficial significant effect on HU of post-molt laying hens' eggs. Of course, in another experiment, organic or inorganic chromium did not affect HU, egg and yolk weight (Eseceli, et al., 2010; Piva et al., 2003). Shell thickness was influenced by the level of 800 ppb N-Cr chloride than control (0 ppb N-CR chloride). In other experiment indicated that supplemental Cr in diet had increasing effects on quail egg shell thickness (Derya et al., 2009; Sahin et al., 2002; Sahin et al., 2001). But, based on investigations of Sirirat et al., (2013), Huseyin et al., (2010) and Yildiz et al., (2004) different supplemental Cr forms did not affect laying quail egg shell thickness.

Chromium is presence in most of tissues (Anderson, 1987). In our experiment, there were not significant effects of N-Cr chloride on blood plasma chromium content among treatments. However, quails in treatment 5 (800 ppb N-Cr chloride) had higher Cr concentration in their blood plasma. This finding is not similar to the investigation of Sirirat et al., (2013). They reported increasing levels of nano chromium picolinate resulted in further retention of Cr in liver, egg yolk and egg shell.

5. Conclusion

In conclusion, it seems that N-Cr level of 800 ppb had further effects on Japanese quail egg internal quality. But there was no effect on Cr content of blood plasma.

Acknowledgments

This article is a part of M.Sc. dissertation in Animal Nutrition, Islamic Azad University, Sanandaj Branch, Kurdistan, IRAN (Thesis supervisor: Dr. Mohammad Amiri Andi).

References

- Anderson, R.A., 1987. Chromium in animal tissues and fluids in: Mertz W. (Ed.). Trace. Elem. Hum. Anim. Nut., Academic Press, New York.
- Anderson, R.A., Bryden, N.A., Evockclover, C.M., Steele, N.C., 1997. Beneficial effects of chromium on glucose and lipid variables in control and somatotropin-treated pigs are associated with increased tissue chromium and altered tissue copper, iron and Zinc. *J. Anim. Sci.*, 75, 657-661.
- Derya, Y., Mustafa, E., 2009. Effects of dietary organic and inorganic chromium supplementation on performance, egg shell quality and serum parameters in pharaoh quails. *J. Biol. Environ. Sci.*, 3, 31-35.
- Eren, M., Baspnar, N., 2004. Effect of dietary CrCl₃ supplementation on some serum biochemical markers in broilers. Influence of season, age and sex. *Revue. de Med. Vet.*, 155, 637-641.
- Eseceli, H., Degirmencioglu, N., Bilgic, M., 2010. The effect of inclusion of chromium yeast (Co-Factor II, Alltech Inc.) and folic acid to the rations of laying hens on performance, egg quality, egg yolk cholesterol, folic acid and chromium levels. *J. Anim. Vet. Adv.* 9, 384-391.
- Holdsworth, E.S., Neville, E., 1990. Effects of extracts of high and low chromium brewer's yeast on metabolism of glucose by hepatocytes from rats fed on high-or lower diets. *Br. J. Nut.*, 63, 623-628.
- Hossain, S.M., Barreto, S.L., Silva, C.G., 1998. Growth performance and carcass composition of broilers fed supplemental chromium from chromium yeast. *Anim. Feed. Sci. Technol.*, 71, 217-228.
- Huseyin, E., Nurcan, D., Murat, b., 2010. The Effect of inclusion of chromium yeast(co-fator II Alltechinc) and folic Acid to the Rations of Laying Hens on performance, Egg quality, egg yolk cholesterol, folic acid and chromium levels. *J. Anim. Vet. Adv.*, 9, 384-391.
- Jensen, L., Chang, S., Wilson, C.H., 1978. Interior egg quality: improvement by distillers' feeds and traces elements. *Poult. Sci.*, 57, 648-654.
- Kim, J.D., Han, I.K., Chae, B.J., Lee, J.H., Park, J.H., Yang, C.J., 1997. Effects of dietary chromium picolinate on performance egg quality, serum traits and mortality rate of brown layers. *Asi. Aust. J. Anim. Sci.*, 10, 1-7.
- Lukaski, H.C., 1999. Chromium as a supplement. *Ann. Rev. Nut.* 19, 279-302.
- NRC, 1994. Nutrient requirements of poultry. 9th Rev. Edn., National Academy Press, Washington, DC., USA.

- Piva, A., Meola, E., Gatta, P.P., Biagi, G.C., Mordenti, A.L., Luchansky, J.B., Silva, S., Mordenti, A., 2003. The effect of dietary supplementation with trivalent chromium on production performance of laying hens and the chromium content in the yolk. *Anim. Feed. Sci. Tech.*, 106, 149-163.
- Sahin, K., Kucuk, O., Sahin, N., 2001. Effects of dietary chromium picolinate supplementation on performance, insulin and corticosterone in laying hens under low ambient temperature. *J. Anim. Physiol. Anim. Nut.*, 85, 142-147.
- Sahin, K., Kucuk, O., Sahin, N., Ozbey, O., 2001. Effects of dietary chromium picolinate supplementation on egg production, egg quality and serum concentrations of insulin, corticosterone, and some metabolites of Japanese quails. *Nut. Res.*, 21, 1315-1321.
- Sahin, K., Ozbey, O., Onderci, M., Cikim, G., Aysondu, M.H., 2002. Chromium supplementation can alleviate negative effects of heat stress on egg production, egg quality and some serum metabolites of laying Japanese quail. *J. Nut.*, 132, 1265-1268.
- Sahin, N., Sahin, K., Onderci, M., Gursu, M.F., Cikim, G., Vijaya, J., Kucuk, O., 2005. Chromium picolinate, rather than biotin, alleviates performance and metabolic parameters in heat-stressed quail. *Br. Poult. Sci.*, 46, 457-463.
- SAS software, 2001. *SAS User's Guide: Statistics, Version 9.2*, SAS Institute, North Carolina, USA.
- Sirirat, N., Lu, J., Hung, A.T., Lien, T., 2013. Effect of different levels of Nanoparticles chromium picolinate supplementation on performance, egg quality, mineral retention and tissues minerals accumulation in layer chicken. *J. Agr. Sci.*, 5, 150-159.
- Uyanik, F., kaya, S., kolsuz, A.H., Eren, M., 2002. The effect chromium Supplementation on egg production, egg quality and serum parameters in laying hens. *Turk. J. Vet. Sci.*, 26, 379-387.
- Yildiz, A.O., Parlat, S.S., Yazgan, O., 2004. The effects of organic chromium supplementation on production traits and some serum parameters of laying quails. *Rev. Med. Vet.* 155, 642-646.
- Zha, L.Y., Xu, Z.R., Wang, M.Q., Gu, L.Y., 2008. Chromium nanoparticle exhibits higher absorption efficiency than chromium picolinate and chromium chloride in Caco-2 cell monolayers. *J. Anim. Physiol. Anim. Nut.*, 92, 131-140.

How to cite this article: Amiri Andi, M., Shahamat, A., 2015. Effects of different levels of nano chromium chloride in diet on egg quality and blood chromium content of layingjapanese quail. *International journal of Advanced Biological and Biomedical Research*, 3(4), 378-383.

Submit your next manuscript to CASRP Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.casrp.co.uk/journals