



Hypercholesterolemia Ameliorating Effect of Bangladeshi High Yield Variety Rice Bran Oil

Mohammad Azizur Rahman^{1*}, Jannatul Ferdous¹, Shakir Hosen², Mahmudul Hasan¹, Afroza Parvin¹, Habibul Bari Shozib²

¹Department of Biochemistry and Molecular Biology, Jahangirnagar University, Savar, Dhaka 1342, Bangladesh

²Grain Quality and Nutrition (GQN) Division, Bangladesh Rice Research Institute (BRRI), Gazipur, Bangladesh

*Corresponding Author E-mail: azizbmb@juniv.edu

Received: 18 July 2020, Revised: 20 August 2020, Accepted: 31 August 2020

ABSTRACT

Background: Hypercholesterolemia led cardiovascular disease (CVD) complications are plaguing the globe. In the present study, Bangladesh Rice Research Institute (BRRI) high yield variety (HYV) rice named as BR5 was selected for extracting bran oil followed by fatty acid analysis and *in vivo* hypocholesterolemic effect of the extracted rice bran oil (RBO) for the sake of ameliorating CVD.

Method: RBO was extracted using n-Hexane followed by fatty acid analysis and anti-hypocholesterolemic study on rats against soybean oil (SBO), mustard oil (MBO) and butter oil (BTO) followed by organ function tests. Study period was four weeks. Then the rats were killed, their blood was collected and serum biochemical parameters were analyzed using semi-auto analyzer.

Results: Fatty acid analyses revealed BR5 was the best among the six HYVs of rice. Compared with the other edible oils, RBO derived from BR5 showed significantly increasing effect upon plasma total cholesterol and low density lipoprotein cholesterol lowering while high density lipoprotein cholesterol increasing. Toxicological studies showed no detrimental effect of BR5 derived RBO on cardiac, renal and hepatic functions.

Conclusion: BR5 is a rich source of RBO that provides balanced fatty acid support and exerts hypocholesterolemic effects without any detectable side effects. Consumption of BR5 derived RBO would aid in CVD amelioration.

Key words: Cardiovascular diseases, Fatty acid profile, High yield variety, Hypercholesterolemia, Rice bran oil, Toxicological studies

1. Introduction

Cardiovascular diseases (CVD) stand among the leading causes of death globally (WHO, CVD FACT SHEET, 2020). Current dietary pattern and sedentary lifestyle have been responsible for

accruing CVD [1, 2]. Thus, CVD management recommends dietary modification. Among them, amelioration of hypercholesterolemia and attenuation of oxidative stress (OS) seem apt. Multiple strategies have been formulated

worldwide in managing hypercholesterolemia and OS. Though therapeutic approaches have shown positive outcomes, they are expensive, render side effects and not easy to reach to the common mass, especially of the developing nation [3, 4]. Besides, consumers' interest in natural, food-based items seems higher than those of the chemically synthesized drugs. Thus, search for cost-effective, easily available, safe and food-based CVD modifying agents has become fashionable. In this context, edible oils of different origin have attracted scientific notion [5, 6].

Rice bran oil (RBO) has been gaining popularity in different cuisine over the last few years. RBO has been reported to possess relatively higher antioxidants and balanced fatty acid composition [7-9]. RBO has been hailed over other edible oils owing to its richer content of bio-components such as antioxidants: γ -oryzanol, tocopherols and tocotrienols [7-9]. So far, γ -oryzanol has been regarded as the most active bio-component providing multitude of health promoting effects [7-9].

Though studies in different countries on non-human primates and on humans have shown promising effect of RBO in ameliorating CVD complications, research in the context of Bangladesh is little researched [5, 6]. Thus, the current experiment was designed to evaluate the efficacy of RBO derived from Bangladeshi rice (DHAAN, in Bengali), in ameliorating experimentally induced hypercholesterolemia in animal models.

Rice is the staple food in Bangladesh and accounts for 80% of the cropped area (KNOWLEDGE BANK BRRI, 2020). 3 million tons of rice bran (at 10 percent of the weight of clean rice) could be produced from about 34.75 million tons of clean rice in Bangladesh annually (KNOWLEDGE BANK BRRI, 2020). Total estimated bran production by the auto and semi-auto rice mill is about 0.9

million ton which could produce 0.20 million ton rice bran oil [10].

Bangladesh consumes about 1.3 million tons of edible oil a year mostly imported by private refiners. Bangladesh faces shortage of in edible oil. Locally produced oil can hardly meet 30 percent of the country's total requirement. Rice bran oil produced from rice mills could substitute for a huge quantity of the imported edible oil every year.

Fulfilling the country's demand of edible oil, quality RBO could be exported and boost national economy. Quality of RBO would be determined based on their capability on health promotion and disease modification. Thus, the present study aimed at comparing the fatty acid content of the selected [previously selected based on higher antioxidative potentialities, data not shown in this study] high yield varieties (HYV) of rice (named as BR5, BR 11, BR 28, BR29, BR48 and BR 49) developed by Bangladesh rice research institute (BRRI) followed by extraction of RBO from the best variety and determination of the best variety's effect on lipid profile modification along with its toxicological studies on rats.

2. Experimental

2.1. Selection and extraction of RBO

Among six HYV developed by BRRI, BR5 was chosen based on *in vitro* antioxidative potentiality [data not shown in this article] for RBO extraction. RBO was extracted using *n*-Hexane followed by de-gumming, de-waxing, bleaching and de-deodorizing. Fatty acid profiling of RBO was followed by determination of its antihypercholesterolemic effect on rat models against soybean oil (SBO), mustard oil (MBO) and butter oil (BTO) as standards.

2.2. Fatty acid analysis

Fatty acid analyses of the selected RBO were performed following the modified

method [11]. Gas chromatographic (GC) analysis of fatty acid methyl esters were carried out using a Nucon 5700 gas chromatograph (India). Stainless steel 10% Silar 7C column packed with 60-120 mesh Gas Chrom Q was used. Maintaining both injector and detector temperature at 240 °C, column temperature was initially set at 160 °C for 5 min followed by ramping a rate of 5 °C per min to the final temperature of 220 °C for 20 minutes. Identification of individual fatty acids had been performed based on the received peak area followed by the quantification against external calibration curve.

2.3. In vivo experiments

Twenty five male Long-Evans rats (12 weeks old), weighing 150 ± 2 g were used in this study. Animals were housed individually in cages in a room at animal house of Bangladesh Rice Research Institute (BRRI). Animals were maintained at 22–24 °C with a controlled 12 hrs light–dark cycle and had free access to tap water and commercial rat feed. Rats were randomly divided into five groups namely control, RBO, SBO, MTO, BTO, each group containing five rats. Each rat was given 15 g feed per day along with 10 g respective oil per rat per day for 28 days.

Rats were anaesthetized using diethyl ether and blood collected from left jugular vein. Blood samples were centrifuged at 6000 rpm for 15 minutes, serum collected

and stored at 4 °C in a refrigerator. Biochemical parameters including liver function tests such as Cholesterol, TG, AST, ALT, ALP, Kidney function tests such as Urea, Creatinine, Uric Acid, Total protein, Albumin, A:G ratio and thyroid hormones such as TSH, FT4 were measured for all groups using commercially available reagent kits following manufacturer's instructions and a semi-auto biochemical analyzer. All the experimental procedure were approved by the ethical committee of BRRI.

2.4. Statistical analyses

Data of triplicate studies were presented as mean \pm SEM. One-way ANOVA and Duncan's multiple range test (DMRT) were computed using SPSS version 20.0.

3. Results and discussion

3.1. Fatty acid profiling

Fatty acid profiling of popular high yielding rice varieties (named as BR5, BR 11, BR 28, BR29, BR48 and BR 49) indicate the range of saturated fatty acids from 16.59 to 21.97 %, mono unsaturated fatty acids from 39.82 to 49.95% and unsaturated fatty acids from 78.03 to 83.88%. RBO possesses the highest level of monounsaturated fatty acid (MUFA), oleic Acid (C18:1) (48.03%) (Tables 1, 2). Our findings are in close proximity with previous research [12].

Table 1. Fatty acid profiling of popular high yielding rice varieties in Bangladesh

Fatty acid composition of rice bran oils					
Fatty acid %	BR11	BRRI dhan28	BRRI dhan29	BRRI dhan48	BRRI dhan49
Lauric acid	ND	0.66	ND	ND	ND
Myristic acid	0.24	0.61	ND	0.44	1.00
Palmitic acid	16.43	18.39	14.10	16.74	17.44
Stearic acid	1.52	1.78	1.74	1.91	2.18
Arachidic acid	0.70	0.54	0.75	0.83	0.97
Behenic acid	0.33	ND	ND	0.38	ND
SFA%	19.21	21.97	16.59	20.29	21.60
Oleic acid	43.16	44.79	49.95	45.36	39.39
Eicosenoic acid	0.44	ND	ND	0.4637	0.4356
MUFA%	43.60	44.79	49.95	45.82	39.82
Linoleic acid	35.98	32.13	32.58	32.66	37.26
Linolenic acid	1.21	1.11	0.88	1.22	1.32
PUFA%	37.19	33.24	33.46	33.88	38.58
UFA%	80.79	78.03	83.41	79.71	78.40

Table 2. Fatty acid profiling of BR5 derived RBO and standard edible oils

Fatty acid profiling of edible oils	RBO (BR5)	MTO	SBO	BTO
SFA%	16.62	15	12	22
a) Myristic acid (C14:0)	00.22	ND	00.5	ND
b) Palmitic acid (C16:0)	12.95	96	11	35
c) Stearic acid (C18:0)	3.02	00.4	3.5	10
d) Arachidic acid (C20:0)	00.43	ND	ND	ND
e) Behenic (C22:0)	ND	2	ND	ND
2. UFA (%)	83.88	88	85	23
i) MUFA (%)	48.32	60	37	35
a) Oleic acid (C18:1)	48.03	13	37	20
b) Eicosenoic acid (C20:1)	00.29	ND	ND	ND
c) Erucic acid (C22:0)	ND	47	ND	ND
ii) PUFA (%)	35.06	28	48	25
a) Linoleic acid (C18:2)	33.98	21.2	43	23
a) Linoleic acid (C1 Feed 8:3)	01.07	6.8	05.0	1.9

3.2. Effect of RBO feeding on body weight change of the rats

During the four-week long observation, gradual increase in body weight (ranging from 140 ± 5 g up to 218 ± 2 g) of the rats of all the groups was noticed (Figure 1). However, there was distinct body weight

change among different groups. The BTO group gained the maximum weight and their rate of becoming weighty surpassed that of the others. Compared with other groups, increased body weight of the BTO group was statistically significant ($P < 0.05$).




Figure 1. Body weight change of the rats fed with RBO and standard edible oils. Where, FEED=commercially available rat food pellet; RBO=rice bran oil; SBO=soybean oil; MTO=mustard oil; BTO=butter oil

3.3. Biochemical parameters

3.3.1. Effect of RBO feeding on lipid profile of the rats

Antihypercholesterolemic and other health promoting effect of RBO were evaluated comparing with those of SBO, MTO and BTO (Table 3). Obtained data

reveals that consuming RBO for 28 days caused significant reduction of plasma total cholesterol level than those of SBO, MTO and BTO fed animals. Compared with the SBO, MTO and BTO fed animals; there was significant reduction in the plasma total cholesterol (TC) level of the RBO fed animals. Though there was a little increase in TC level of the RBO group than those of the normocholesterolemic control (feed)

group, this increment was statistically insignificant (Table 3). On the contrary, all the SBO, MTO and BTO fed animals significantly increased level of plasma TC levels than those of the control group. Similar trend was observed in case of plasma triacylglycerol levels. RBO showed plasma HDL-C increased while LDL-C and VLDL-C effects decreased (Table 3). However, this effect was not uniformly significant in all groups (Table 3).

Our findings are in concordance with those of [13]. Poly unsaturated fatty acid (PUFA, 35.06%) composition, especially linoleic acid (33.98%), along with non-fatty acid components such as the micronutrients (triterpene alcohols, phytosterols, γ -oryzanol and tocotrienols) present in the unsaponifiable fraction of RBO might had synergistically lowered rat serum cholesterol level [13]. Hypocholesterolemic effect of RBO might also be attributed to fecal excretion of cholesterol [14, 15].

Triacylglycerol (TG) level increased insignificantly in the RBO fed animals, significantly while in the rest. Earlier studies on rats indicate that rats fed diets containing 10% RBO for 8 weeks lowered plasma TC, LDL-C and VLDL-C and increased HDL-C levels, while no changes in TG level on cholesterol-containing or cholesterol-free diets were observed [15].

RBO caused significant increment in plasma HDL-C level and decrement in LDL-C and VLDL-C levels. Similar trend was observed for SBO and MTO while BTO had opposite effects. The reduction of TG and TC associated with LDL-C suggest that cholesterol lowering activity of RBO might result from the enhanced catabolism of LDL-C through hepatic receptors for final elimination in the form of bile acids [16]. There exists a positive correlation between an increased LDL-C/HDL-C ratio and the development of atherosclerosis.

Increased level of HDL-C in the RBO fed rats indicates its potent protective action against atherogenesis since an independent inverse relationship exists between plasma HDL-C levels and cardiovascular risk. Underlying mechanism may be attributed to the enhancement of lecithin cholesteryl acyl transferase (LCAT), playing a key role in incorporating the free cholesterol into HDL and transferring back to VLDL or intermediate density lipoproteins (IDL), which is taken back by the liver cells [17]. Additional mechanism may involve an inhibition of hepatic triglyceride lipase (HTL) on HDL which may lead to a rapid catabolism of blood lipids through extrahepatic tissues [18]. Thus, the current findings confirm the anti-hyperlipidaemic properties of BR5 derived RBO and its overall impact to lower the CVD risk.

Table 3. Plasma lipid profile of rats

Treatments	Total cholesterol (mg/dl)	TG (mg/dl)	HDL-C (mg/dl)	LDL-C (mg/dl)	VLDL-C (mg/dl)
Feed	45.68± 0.12 ^a	61.81± 0.48 ^a	29.06 ^a ± 0.42 ^a	36.97 ± 0.58 ^a	24.54 ± 0.13 ^a
RBO+ Feed	47.73± 0.16 ^a	64.36± 0.34 ^a	36.47 ± 0.55 ^b	35.44 ± 0.58 ^a	20.78 ± 0.17 ^b
SBO+ Feed	58.79± 0.22 ^b	92.39± 0.44 ^b	31.94 ± 0.58 ^a	55.97 ± 0.58 ^b	34.40 ± 0.18 ^c
MTO+ Feed	67.71± 0.27 ^c	109.41± 0.54 ^c	30.33 ± 0.40 ^a	66.97 ± 0.58 ^c	33.47 ± 0.18 ^d
BTO+ Feed	86.23± 0.29 ^d	161.16± 0.52 ^d	28.73 ± 0.40 ^a	70.97 ± 0.58 ^d	36.67 ± 0.10 ^e

Results have been expressed as mean±SEM of triplicate determination. The number of rats for each group was 5. Data in the same column having different superscripts are significantly different at $P < 0.05$, as measured by the DMRT Test.

3.3.2. Cardiac and liver function tests

Enzymatic tests were performed to assess the safety issue of RBO and other edible oils on the experimental subjects. Compared with the control group, we observed significantly increased levels of serum AST/GOT, ALT/GPT and ALP

enzymes ($P \leq 0.05$) were observed in the SBO, MTO and BTO rats; however, there was little and insignificant increment in case of RBO rats (Table 4). The obtained results suggest that compared with SBO, MTO and BTO, the RBO had less toxic effect on rats' cardiac and hepatic tissues. Our findings are in concordance with previous research [13, 19]. Thus, BR5 derived RBO had no detectable detrimental effect on cardiac and renal functions. Also, no rat died, nor did they show behavioral abnormalities during the experimental period.

Table 4. Cardiac and liver function tests

Treatments	AST (U/L)	ALT (U/L)	ALP (U/L)
Feed	229.27± 0.59 ^a	504.12± 0.67 ^a	35.33± 0.34 ^a
RBO+ Feed	233.16± 0.64 ^a	506.16± 0.45 ^a	37.11± 0.30 ^a
SBO+ Feed	300.37± 0.44 ^b	514.10± 0.18 ^a	45.33± 0.48 ^b
MTO+ Feed	325.36± 0.74 ^c	528.17± 0.33 ^a	57.23± 0.23 ^c
BTO+ Feed	357.66± 0.29 ^d	533.13± 0.39 ^a	82.17± 0.27 ^d

Results have been expressed as mean±SEM of triplicate determination. The number of rats for each group was 5. The data in the same column having different superscripts are significantly different at $P < 0.05$, as measured by the DMRT Test.

3.3.3. Kidney function tests

Kidney function parameters, i.e. creatinine, urea, and uric acid, indicate that

RBO had very little increasing effect on plasma levels of creatinine, urea and uric acid (Table 5). However, SBO, MTO and BTO exerted comparatively higher effects. Our findings are in concordance with previous research [13]. Thus, from the toxicological point of view, BR5 derived RBO could be considered as safe.

Table 5. Kidney function tests

Treatments	Creatinine (mg/dl)	Urea (mg/dl)	Uric acid (mg/dl)
Feed	0.52± 0.02 ^a	35.1± 0.29 ^a	1.48± 0.12 ^a
RBO+ Feed	0.53± 0.05 ^a	37.66± 0.22 ^a	1.61± 0.14 ^a
SBO+ Feed	0.59± 0.09 ^b	44.33± 0.19 ^b	2.23± 0.09 ^b
MTO+ Feed	0.58± 0.04 ^b	46.66± 0.12 ^b	2.39± 0.11 ^b
BTO+ Feed	0.65± 0.07 ^b	47.66± 0.26 ^b	2.80± 0.20 ^b

Results have been expressed as mean±SEM of triplicate determination. The number of rats for each group was 5. The

data in the same column having different superscripts are significantly different at $P < 0.05$, as measured by the DMRT Test.

3.3.4. Protein and hormonal levels

Plasma levels of total protein, globulin, albumin/globulin ratio, free T4 (FT4; thyroxine) and thyroid stimulating hormone (TSH) were measured in all the experimental animals (Table 6). There was discrepancy among the groups regarding each of the parameters;

however, this was statistically insignificant. Previous studies suggest TSH lowering effect of γ -oryzanol extracted from rice bran oil [20]. Thus, our findings suggest no detrimental effect of BR5 derived RBO on blood biochemistry profile of the experimental animals.

Table 6. Plasma protein and hormone level

Treatments	Total Protein (g/dl)	Globulin (g/dl)	A/G Ratio	FT4 (mg/dL)	TSH (mIU/L)
Feed	7.15± 0.15 ^a	2.11± 0.13 ^a	2.40± 0.25 ^a	1.46± 0.08 ^a	<0.06
RBO+ Feed	6.66± 0.11 ^a	1.65± 0.05 ^a	3.07± 0.09 ^a	1.26± 0.11 ^a	<0.06
SBO+ Feed	7.00± 0.05 ^a	1.37± 0.09 ^a	4.12± 0.25 ^a	1.47± 0.07 ^a	<0.06
MTO+ Feed	6.89± 0.09 ^a	1.66± 0.05 ^a	3.15± 0.30 ^a	1.82± 0.08 ^a	<0.06
BTO+ Feed	7.42± 0.06 ^a	2.20± 0.08 ^a	2.36± 0.15 ^a	1.93± 0.03 ^a	<0.06

Results have been expressed as mean±SEM of triplicate determination. The number of rats for each group was 5. The data in the same column having different superscripts are significantly different at $P < 0.05$, as measured by the DMRT Test.

4. Conclusions

Plasma cholesterol lowering effect of RBO derived from the high yield rice variety BR5 surpasses those of SBO, MTO and BTO. Toxicological studies suggest this RBO to be safe. Thus, BR5 derived RBO could replace the other edible oils usually utilized in our daily cooking and dressing. This would aid in modification of current dietary pattern towards healthy life style practicing for the sake of overcoming CVD complications. However, large scale studies of BR5 derived RBO on human sample are warranted. Meanwhile, in countries, where production cost of RBO is higher, combination of RBO with low cost edible oils could be considered.

Abbreviations

A:G ratio= Albumin/Globulin ratio
ALP= Alanine phosphatase

ALT= Alanine amino transferase
ANOVA= Analysis of variance
AST= Aspartate amino transferase
BRRI= Bangladesh rice research institute
BTO= Butter oil
CVD= Cardiovascular diseases
GC= Gas chromatographic analysis
HDL-C= High density lipoprotein-cholesterol
LDL-C= Low density lipoprotein-cholesterol
VLDL-C= Very low-density lipoprotein cholesterol
MTO= Mustard oil
MUFA= Monounsaturated fatty acid
PUFA= Polyunsaturated fatty acid
RBO= Rice bran oil
SBO= Soybean oil

Conflict of interest

All authors declare that they have no conflict of interest/All authors declare no conflict of interest.

Consent for publications

All authors have read and approve the final manuscript for publication/We have read and approve the final manuscript for publication.

Availability of data and material

We have embedded all data in the manuscript.

Authors' contributions

Study design, supervision, analysis of data and manuscript writing had been performed by Mohammad Azizur Rahman and Habibul Bari Shozjib. Animal handling and data collection had been performed by Jannatul Ferdous and Shakir Hossen. Mahmudul Hasan and Afroza Parvin edited the manuscript.

Ethics approval and consent to participate

Prior approval from animal experimentation ethics committee of Bangladesh Rice Research Institute had been collected.

Acknowledgement

Authors gratefully acknowledge the Bangladesh Rice Research Institute (BRRI) and Jahangirnagar University authorities for providing support in respect of animal housing, laboratory tests and necessary grant-in-aid to conduct this research.

5. References

- Pan XF, Li Y, Franco OH, Yuan JM, Pan A, Koh WP. (2019). Impact of combined lifestyle factors on all-cause and cause-specific mortality and life expectancy in Chinese: the Singapore Chinese Health Study. *J. Gerontol. A Biol. Sci. Med. Sci.*, <https://doi.org/10.1093/gerona/glz271>
- Hansen D, Niebauer J, Cornelissen V, Barna O, Neunhäuserer D, Stettler C, Tonoli C, Greco E, Fagard R, Coninx K, Vanhees L, Piepoli MF, Pedretti R, Ruiz G R, Corrà U, Schmid JP, Davos C H, Edelmann F, Abreu A, Rauch B, Ambrosetti M, Braga SS, Beckers P, Bussotti M, Faggiano P, Garcia-Porrero E, Kouidi E, Lamotte M, Reibis R, Spruit MA, Takken T, Vigorito C, Völler H, Doherty P, Dendale P. (2018). Exercise Prescription in Patients with Different Combinations of Cardiovascular Disease Risk Factors: A Consensus Statement from the EXPERT Working Group. *Sports Med*, 48(8): 1781-1797. <https://doi.org/10.1007/s40279-018-0930-4>
- Hartley L, Mavrodaris A, Flowers N, Ernst E, Rees K. (2017). WITHDRAWN: Transcendental meditation for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev*, 11(11): CD010359. <https://doi.org/10.1002/14651858.CD010359.pub3>
- Davari A, Solouki M, Fazeli-Nasab B. (2018). Effects of jasmonic acid and titanium dioxide nanoparticles on process of changes of phytochemical and antioxidant in genotypes of *Satureja hortensis* L. *Eco-Phytochemical Journal of Medicinal Plants*, 5(4): 1-20 (In Persian).
- Sharifi-Rad J, Rodrigues CF, Sharopov F, Docea AO, Can Karaca A, Sharifi-Rad M, Kahveci Karıncaoglu D, Gülseren G, Şenol E, Demircan E, Taheri Y, Suleria HAR, Özçelik B, Nur Kasapoğlu K, Gültekin-Özgülven M, Daşkaya-Dikmen C, Cho WC, Martins N, Calina D. (2020). Diet, Lifestyle and Cardiovascular Diseases: Linking Pathophysiology to Cardioprotective Effects of Natural Bioactive Compounds. *Int J Environ Res Public Health*, 17(7). <https://doi.org/10.3390/ijerph17072326>
- Salehi B, Del Prado-Audelo ML, Cortés H, Leyva-Gómez G, Stojanović-Radić Z, Singh YD, Patra JK, Das G, Martins N,

- Martorell M, Sharifi-Rad M, Cho WC, Sharifi-Rad J. (2020). Therapeutic Applications of Curcumin Nanomedicine Formulations in Cardiovascular Diseases. *J Clin Med*, 9(3).
<https://doi.org/10.3390/jcm9030746>
7. Ali MA, Islam MA, Othman NH, Noor AM, Hossen J, Ibrahim M. (2019). Effect of Heating on Compositional Characteristics and Oxidative Stability of Crude and Refined Rice Bran Oil. *J Oleo Sci*, 68(11): 1085-1097.
<https://doi.org/10.5650/jos.ess19140>
8. Kobayashi E, Ito J, Kato S, Sawada K, Matsuki M, Hashimoto H, Miyazawa T, Nakagawa K. (2016). Presence of orally administered rice bran oil γ -oryzanol in its intact form in mouse plasma. *Food Funct*, 7(12): 4816-4822.
<https://doi.org/10.1039/c6fo01552b>
9. Sohail M, Rakha A, Butt M S, Iqbal M J, Rashid S. (2017). Rice bran nutraceuticals: A comprehensive review. *Crit Rev Food Sci Nutr*, 57(17): 3771-3780.
<https://doi.org/10.1080/10408398.2016.1164120>
10. Rahman M, Zaman M, Islam A S. (2011). Prospect of rice bran oil production in Bangladesh. *J Agril Engg*, 39(1): 1-8.
11. Deshpande A B, Chidley H G, Oak P S, Pujari K H, Giri A P, Gupta V S. (2016). Data on changes in the fatty acid composition during fruit development and ripening of three mango cultivars (Alphonso, Pairi and Kent) varying in lactone content. *Data in brief*, 9: 480-491.
<https://doi.org/10.1016/j.dib.2016.09.018>
12. Sugano M, Tsuji E. (1997). Rice bran oil and cholesterol metabolism. *J Nutr*, 127(3): 521S-524S.
<https://doi.org/10.1093/jn/127.3.521S>
13. Al-Okbi SY, Mohamed DA, Hamed TE, Al-Siedy ES. (2019). Rice bran as source of nutraceuticals for management of cardiovascular diseases, cardio-renal syndrome and hepatic cancer. *Journal of Herbmed Pharmacology*, 9(1): 68-74.
<https://doi.org/10.15171/jhp.2020.10>
14. Lei L, Chen J, Liu Y, Wang L, Zhao G, Chen Z Y. (2018). Dietary Wheat Bran Oil Is Equally as Effective as Rice Bran Oil in Reducing Plasma Cholesterol. *J Agric Food Chem*, 66(11): 2765-2774.
<https://doi.org/10.1021/acs.jafc.7b06093>
15. Sunitha T, Manorama R, Rukmini C. (1997). Lipid profile of rats fed blends of rice bran oil in combination with sunflower and safflower oil. *Plant Foods Hum Nutr*, 51(3): 219-230.
<https://doi.org/10.1023/a:1007993016796>
16. Khanna A K, Rizvi F, Chander R. (2002). Lipid lowering activity of *Phyllanthus niruri* in hyperlipemic rats. *J Ethnopharmacol*, 82(1): 19-22.
[https://doi.org/10.1016/s0378-8741\(02\)00136-8](https://doi.org/10.1016/s0378-8741(02)00136-8)
17. Devi R, Sharma D K. (2004). Hypolipidemic effect of different extracts of *Clerodendron colebrookianum* Walp in normal and high-fat diet fed rats. *J Ethnopharmacol*, 90(1): 63-68.
<https://doi.org/10.1016/j.jep.2003.09.022>
18. Anila L, Vijayalakshmi N R. (2002). Flavonoids from *Embllica officinalis* and *Mangifera indica*-effectiveness for dyslipidemia. *J Ethnopharmacol*, 79(1): 81-87. 10.1016/s0378-8741(01)00361-0
19. Araghi A, Seifi S, Sayrafi R, Sadighara P. (2016). Safety assessment of rice

bran oil in a chicken embryo model.

Avicenna J Phytomed, 6(3): 351-356.

20. Shimomura Y, Kobayashi I, Maruto S, Ohshima K, Mori M, Kamio N, Fukuda H. (1980). Effect of gamma-oryzanol

on serum TSH concentrations in primary hypothyroidism. *Endocrinol Jpn*, 27(1): 83-86.

<https://doi.org/10.1507/endocrj1954.27.83>

How to cite this article: Mohammad Azizur Rahman, Jannatul Ferdous, Shakir Hosen, Mahmudul Hasan, Afroza Parvin, Habibul Bari Shozib, Hypercholesterolemia Ameliorating Effect of Bangladeshi High Yield Variety Rice Bran Oil. *International Journal of Advanced Biological and Biomedical Research*, 2021, 9(1), 48-57. Link: <http://www.ijabbr.com/article 44619.html>