



IJABBR- 2014- eISSN: 2322-4827

International Journal of Advanced Biological and Biomedical Research

Journal homepage: www.ijabbr.com



Review Article

Physiology of Plants Affected by Ageing

Morad Shaban¹ and Zahra Rahmati Motlagh^{1*}

¹Young Researchers and Elite club, Boroujed branch, Islamic Azad University, Boroujed, Iran

ARTICLE INFO

Article history:

Received: 14 May, 2014

Revised: 29 May, 2014

Accepted: 2 July, 2014

ePublished: 30 July, 2014

Key words:

Ageing

Physiology of seeds

Reactive oxygen species

Seed damage

ABSTRACT

Objective: Ageing of the stored seed is a natural phenomenon and the seeds tend to lose viability even under ideal storage conditions. Accelerated ageing has been recognized as a good predictor of the storability of seed lots. Aged seeds show decreased vigor and produce weak seedlings that are unable to survive once reintroduced into a habitat. The characteristics of the chemical composition of oil crops (soybean and sunflower) are related to specific processes occurring in seed during storage. Lipid auto-oxidation and increase of free fatty acid content during storage are the most often mentioned reasons for accelerated damage of seed of oily plant species.

INTRODUCTION

Although highest seed yield in agriculture in normal condition of fertilizing and environmental conditions (Shaban, 2013a, b; Beyranvand et al. 2013; Kiani et al. 2013), but aged seeds has a low yield in best condition same above. During the last 20 years, considerable research has been conducted to understand better the physiology of seed ageing, since the primary processes and their interactions involved in seed ageing are not yet fully understood (McDonald, 1999). During storage, a number of physiological and physicochemical changes occur, termed aging (Sisman, 2005). The rate at which the seed aging process takes place depends on the ability of seed to resist degradation changes and protection mechanisms, which are specific for each plant species (Sisman and Delibas, 2004; Mohammadi et al. 2011). The main external factors causing seed damage during storage are the temperature, relative air humidity and oxygen. Possibility to regulate these factors makes the

basis for longer seed storage. Seed deterioration is inexorable and the best that can be done is to control its rate. Many factors contribute to the predisposition for seed deterioration. These include genetics where certain seeds are inherently longer lived than others. Seed structure can also influence seed deterioration. Simple differences in seed size can mean that smaller seeds with a greater surface area to volume ratio are more exposed to uptake of water that would make them prone to deterioration more than larger seeds. Seed chemistry influences the amount of free water available to seeds that increase deterioration rate. Seeds that possess mucilage around their seed coats such as *Salvia* under high relative humidity environments would be more likely to transfer that moisture into the seed causing more rapid deterioration. It has been known for a long time that a progressive fragmentation of embryonic nuclear DNA occurs during seed ageing (Cheah and Osborne, 1978). DNA damage can be due to an uncontrolled degradation following extensive DNA

*Corresponding Author: Zahra Rahmati Motlagh, Young researchers and elite club, Boroujed branch, Islamic Azad University, Boroujed, Iran (Rahmati.z1365@yahoo.com)

oxidation or to DNA laddering, as is commonly observed in active and genetically controlled programmed cell death (PCD) (Stein and Hansen, 1999). ROS have been widely cited as being the main factor causing seed ageing during their prolonged storage (Priestley, 1986). Deterioration in different tissues and seeds is a composite of tissues that differ in their chemistry and proximity to the external environment. Thus, it should not be assumed that seed deterioration occurs uniformly throughout the seed. Perhaps the best example that this does not occur comes from the use of the tetrazolium chloride (TZ) test where living tissues in a seed turn red when exposed to the compound. When studies have been conducted on seeds using controlled natural and artificial aging conditions, differences in the deterioration of seed tissues have been observed. In dicot seeds such as soybean, the embryonic axis is more sensitive to deterioration than the cotyledons. Embryonic axes possess more MDA and total peroxides which are markers of greater free radical presence in the axes compared to aged cotyledons.

2.1. Ageing of seeds

Inactivation and degradation of enzymes due to changes in their macromolecular structures is one of the most important hypotheses proposed regarding causes of ageing in seeds (Bailly, 2004; Basavarajappa et al. 1991; Basra and Malik, 1994; Goel et al. 2002; Kalpana and Rao, 1993; Lehner et al. 2008; McDonald, 2004; Salama and Pearce, 1993). Most of these studies suggest that decreases occur in the activity of enzymes such as superoxide dismutase, catalase, peroxidase and glutathione reductase in aged seeds. The general decrease in enzyme activity in the seed lowers the respiratory capacity, which in turn lowers both the energy (ATP) and assimilates supply of the germinating seed. Therefore, several changes in the enzyme macromolecular structure may contribute to their lowered germination efficiency. As evidence mounts, the leading candidate causing seed deterioration increasingly appears to be free radical production. Free radical production, primarily initiated by oxygen, has been related to the peroxidation of lipids and other essential compounds found in cells. This causes a host of undesirable events to include decreased lipid content. Lipid peroxidation begins with the generation of a free radical (an atom or a molecule with an unpaired electron) either by autoxidation or enzymatically by oxidative enzymes such as lipoxygenase present in many seeds. One of the major sources of ROS in metabolically active seeds is the mitochondrial respiratory chain (Bailly, 2004). Lipid peroxidation occurs in all cells, but in fully imbibed cells, water acts as a buffer between the autoxidatively-generated free radicals and the target macromolecules, thereby reducing damage. Thus, as seed moisture content is lowered, autoxidation is more common and is accelerated by high temperatures and increased oxygen concentrations. Lipid autoxidation may

be the primary cause of seed deterioration at moisture contents below 6%. Above 14% moisture content, lipid peroxidation may again be stimulated by the activity of hydrolytic oxidative enzymes such as lipoxygenase, becoming more active with increasing water content. Between 6 and 14% moisture content, lipid peroxidation is likely at a minimum because sufficient water is available to serve as a buffer against autoxidatively generated free radical attack, but not enough water is present to activate lipoxygenase-mediated free radical production. The relationship between seed deterioration and the enzymes involved in lipid peroxidation, free radical removal, and seed respiratory metabolism also has been studied (Shen and Odén, 1999). All enzyme activity is positively correlated with germination of seed as ageing progressed germination also decreased and enzyme activity also decreased which showed significant deterioration in both accelerated as well as in natural aged seed lot. All seeds undergo aging process during long-term storage which leads to deterioration in seed quality, especially in the humid tropical regions. However, the rate of seed deterioration can vary among various plant species (Merritt et al. 2003). Aged seeds show decreased vigor and produce weak seedlings that are unable to survive once reintroduced into a habitat (Atici et al. 2007). Some protective mechanisms involving free radical and peroxide scavenging enzymes, such as catalase (CAT), peroxidase (POD) and superoxide dismutase (SOD) have been evaluated within the mechanism of seed aging (Hsu et al. 2003, Goel et al. 2003, Pukacka and Ratajczak 2007). Loycrajjou et al. (2008) reported that ageing induced deterioration increase the extent of protein oxidation thus inducing loss of functional properties of proteins and enzymes.

2.2. Various seeds and ageing

A fatty acid composition is the most important factor which determines oils susceptibility to oxidation (Morello et al. 2004). Quality parameters of seed such as oil content, fatty acid composition and protein content are significantly influenced by storage conditions and time (Ghasemnezhad and Honermeier 2007). For example, sunflower seed storage demands special care due to high oil content which can easily provoke processes that can lead to loss of germination and viability. For short term storage the seed may be kept in storage facilities that are dry and relatively cold, in order to preserve biological value of seed. These facilities have no special temperature and humidity control. For longer and more reliable seed storage, facilities with specific climate control should be used. Depending on temperature coefficient the plant species display different reaction to storage longevity at a constant storage temperature of 10°C. Onion seed showed the lowest temperature coefficient, and it proved to be hard for storing, and the barley seed showed the highest temperature coefficient, and it maintained its germ inability under various conditions (Milošević et al. 1996).

Seed moisture content is the most important determinant of longevity in storage. The important thing from the perspective of chemical reactions in the seed is water activity, which means the chemical potential of water in the system (Basra, 1984). Storage of one kind of seed at 14% moisture may be feasible, while for a different species 14% moisture might be too high. Seed of the majority of agricultural plant species may be stored for several years provided that the seed moisture is maintained between 5-8%. The seed moisture content depends on relative air humidity in the seed storage facility. Higher air humidity increases seed moisture content, leading to more rapid seed deterioration, particularly at the moisture content above 12%. Seed with 70% of germination will generally not store as well as another one with 95% of germination. But it is apparent that some important aspects of seed storability and vigor may not be reflected in the initial germination percent prior to storage. A high initial germination provides no assurance that the lot will store as well as another lot of the same kind with the same or even lower germination. The solution to problems to the storability of seed lies in the development of tests that will differentiate among seed lots with respect to storage potential. Seed vigor declines first as seed deteriorates, followed by loss of germination and viability. In the majority of plant species having oil rich seeds the lipids that are at risk of auto-oxidation of fatty acid chain. Decrease of linoleic acid content in aged seed was highly pronounced in maize lines with higher oil content (Balešević-Tubić et al. 2004). It is believed that biochemical processes of lipid peroxidation are the major cause of seed deterioration during storage. Lipid peroxidation and products resulting from these processes lead to DNA denaturation, prevent translation and protein transcription, and cause oxidation of the most reactive amino acids (Popović et al. 2006). When these types of damages occur in seed, they may cause decrease in vigor and seed germination.

References

Bailly, C., 2004. Reactive oxygen species and antioxidants in seed biology. *Seed Science Research*. 14; 93-107.

Balešević-Tubić, S., Tatić, M., Miladinović, J., Malenčić, Đ., 2004. Lipid peroxidation and activity of superoxide dismutase associated with natural aging of oil maize seed. Abstracts. 27th ISTA Congress Seed Symposium. Budapest, Hungary.

Balešević-Tubić, S., Tatić, M., Miladinović, J., Pucarević, M., 2007. Changes of fatty acids content and vigor of sunflower seed during natural aging. *Helia*. 30; 61-67.

Mechanism of oxidative damage is very complex and occurs as two different types of fatty acid changes. The first one is linked to the process of aging during the first week of storage and includes spontaneous oxidation of unsaturated fatty acids, with no changes occurring in saturated fatty acids. In the second type of damage the seed that has lost its ability to germinate showed oxidation of both saturated and unsaturated fatty acids. Activity of free radicals in seed may depend on water content, seed components (for example the whole seed, cotyledon, embryo), seed lot, species and variety, as well as the aging treatment (Laloi et al. 2004). Peroxidative changes in fatty acid composition of membrane lipids exert influence on viscosity, permeability and membrane cell function. Also, decrease in mitochondrial respiration during storage could be associated with peroxidative changes in lipid mitochondria that lead to loss of seed vigor (Ferguson et al. 1990). Final product of lipid peroxidation is lipid hydroperoxid (ROOH) from which aldehydes are formed, including malonyl-dialdehyde (MDA). Determination of MDA content is the conventional method used for determination of lipid peroxidation (Sung and Jeng, 1994).

Conclusion

In this study we understand that aging in all seeds is a serious problem that induces damage to seed vigor and loss of longevity of seeds. These problems reduce germination of seeds that laid to loss of green surface in agricultural farm and loss of seed yield in all seeds crop. The characteristics of the chemical composition of oil crops are related to specific processes occurring in seed during storage. Lipid auto-oxidation and increase of free fatty acid content during storage are the most often mentioned reasons for accelerated damage of seed of oily plant species.

Basavarajappa, B.S., Shetty, H.S., Prakash, H.S. 1991. Membrane deterioration and other biochemical changes, associated with accelerated ageing of maize seeds. *Seed Science and Technology*. 19; 279-286.

Basra, A.S., Malik, C.P., 1994. Amelioration of the effects of ageing in onion seeds. *Biology Plantarum*. 36(3); 365-371.

Basra, A.S., 1984. Basic mechanisms and agricultural implications. *Food Products Press, England*. 183-200.

Beyranvand, H., Farnia, A., Nakhjavan, S.H., Shaban, M., 2013. Response of yield and yield components of maize (*Zea mays* L.) to different bio fertilizers. *International*

Journal of Advanced Biological and Biomedical Research. 1 (9); 1068-1077.

Cheah, K.S.E., Osborne, D.J., 1978. DNA lesions occur with loss of viability in embryos of ageing rye seeds. *Nature*. 272; 593-599.

Stein, J.C., Hansen, G., 1999. Mannose induces an endonuclease responsible for DNA laddering in plant cells. *Plant Physiology*. 121; 71-80.

Copeland, L.O., Mc Donald, M.B., 1995. Principles of seed science and technology. New York: McMillan. 321.

Ferguson, J.M., TeKrony, D.M., Egli, D.B., 1990. Changes during early soybean seed and axes deterioration: II. Lipids. *Crop Science*. 30; 179-182.

Ghasemnezhad, A., Honermeier, B., 2007. Influence of storage conditions on quality and viability of high and low oleic sunflower seeds. *International Journal of Plant Production*. 3(4); 41-50.

Goel, A., Goel, A.K., Sheoran, I.S., 2003. Changes in oxidative stress enzymes during artificial ageing in cotton (*Gossypium hirsutum* L.) seeds. *Journal of Plant Physiology*. 160; 1093-1100.

Goel, A., Goel, A.K., Sheoran, I.S., 2002. Changes in oxidative stress enzymes during artificial ageing in cotton (*Gossypium hirsutum* L.) seeds. *Journal of Plant Physiology*. 160; 1093-1100.

Hsu, C.C., Chen C.L., Chen, J.J., Sung, J.M., 2003. Accelerated aging enhanced lipid peroxidation in bitter melon seeds and effects of priming and hot water soaking treatments. *Scientia Horticulturae*. 98; 201-12.

Kalpana, R., Rao, M.K.V., 1993. Lowered lipoxygenase activity in seeds of pigeonpea (*Cajanus cajan* L. Mill.) cultivars during accelerated ageing. *Seed Science and Technology*. 21; 269-272.

Kiani, M., Farnia, A., Shaban, M., 2013. Changes of seed yield, seed protein and seed oil in rapeseed (*Brassica napus* L.) under application of different bio fertilizers. *International journal of Advanced Biological and Biomedical Research*. 1(10); 1170-1178.

Kibinza, S., Vinel, D., Côme, D., Bailly, C., Corbineau, F., 2006. Sunflower seed deterioration as related to moisture content during ageing, energy metabolism and active oxygen species scavenging. *Physiologia Plantarum*. 128; 496-506.

Laloi, C., Apel, K., Danon, A., 2004. Reactive oxygen signalling: the latest news. *Current Opinion in Plant Biology*. 7; 323-328.

Lehner, A., Mamadou, N., Poels, P., Côme, D., Bailly, C., Corbineau, F., 2008. Changes in soluble carbohydrates, lipid peroxidation and antioxidant enzyme activities in the embryo during ageing in wheat grains. *Journal of Cereal Science*. 47; 555-565.

Loycrajjou, A., Lovigny, Y., Steven, P.C., Groot, B., Belghazi, M., Job, C., Job, D., 2008. Proteome wide characterization of seed ageing in arabidopsis. A comparison between artificial and natural ageing. *Production Plant Physiology*. 148: 620-41.

McDonald, M.B., 1999. Seed deterioration: physiology, repair and assessment. *Seed Science and Technology*. 27; 177-237.

McDonald, M.B., 2004. Orthodox seed deterioration and its repair, pp. 273-304. In: Handbook of Seed Physiology: Applications to Agriculture, Benech-Arnold, R. L. and R.A. Sanchez (Eds.). *Food Products Press*. New York.

Merritt, D.J., Senaratna, T., Touchell, D.H., Dixon, K.W., Sivasithamparam, K., 2003. Seed ageing of four Western Australian species in relation to storage environment and seed antioxidant activity. *Seed Science Research*. 13; 155-65.

Milošević, M., Ćirović, M., Mihaljev, I., Dokić, P., 1996. Opšte semenarstvo, Feljton, Novi Sad: 148-151.

Mitrovic, A., Ducic, T., Rajlic, I.L., Radotic, K., Zivanovic, B., 2005. Changes in *Chenopodium rubrum* seeds with ageing. *Annals of New York Academy of Science*. 1048; 505-8.

Mohammadi, H., Soltani, A., Sadeghipour, H.R., Zeinali, E., 2011. Effect of seed aging on subsequent seed reserve utilization and seedling growth in soybean. *International Journal of Plant Product*. 5(1); 65-70.

Morello, J.R., Motilva, M.J., Tovar, M.J., Romero, M.P., 2004. Changes in commercial virgin olive oil (CV Arbequina) during storage with special emphasis on the phenolic fraction. *Journal of Food Chemistry*. 85; 357-364.

Murthy, U.M.N., Kumar, P.P., and Sun, W.Q., 2003. Mechanisms of seed ageing under different storage conditions for *Vigna radiata* (L.) Wilczek: Lipid peroxidation, sugar hydrolysis, Maillard reactions and

their relationship to glass state transition. *Journal of Food Chemistry*. 54; 1057-1067.

Pallavi, M., Sudheer, S.K., Dangi, K.S., Reddy, A.V., 2003. Effect of seed ageing on physiological, biochemical and yield attributes in sunflower (*Helianthus Annus L.*) cv. Morden. *Seed Research*. 31 (2); 161-8.

Popović, B., 2006. Uticaj γ - zračenja na antioksidantni sistem odabranih genotipova soje i pojava oksidativnog stresa. Doktorska disertacija, Prirodno-matematički fakultet, Departman za hemiju, Novi Sad.

Priestley, D.A., 1986. Seed Ageing: Implications for Seed Storage and Persistence in the Soil. Cornell University Press, Ithaca, New York.

Pukacka, S., Ratajczak, E., 2007. Age-related biochemical changes during storage of beech (*Fagus sylvatica L.*) seeds. *Seed Science Research*. 17; 45-53.

Salama, A.M., Pearce, R.G., 1993. Ageing of cucumber and onion seeds: Phospholipase d, lipoxygenase activity and changes in phospholipid content. *Seed Research*. 44; 1253-1265.

Scialabba, A., Bellani, L.M., Dell'aquila, A., 2002. Effects of ageing on peroxidase activity and localization in radish (*Raphanus sativus L.*) seeds. *European Journal of Histochemistry*. 46 (4); 351-358.

Shaban, M., 2013a. Application of seed equilibrium moisture curves in agro physics. *International journal of Advanced Biological and Biomedical Research*. 1 (9); 885-898.

Shaban, M. (2013b). Biochemical aspects of protein changes in seed physiology and germination. *International journal of Advanced Biological and Biomedical Research*. 1 (8); 885-898.

Shen, T.Y. Oden, P.C., 1999. Activity of sucrose synthase, soluble acid invertase and fumarase in germinating seeds of Scots pine (*Pinus sylvestris L.*) of different quality. *Seed Science and Technology*. 27 (3); 825-838.

Sisman, C., Delibas, L., 2004. Storing sunflower seed and quality losses during storage. *Journal of Central European Agriculture*. 4; 239-250.