



## The Study on Agroclimatic Conditions of Sunflower in Kurdistan Province by Means of GIS System

Behroz Aminzadeh\*, Sirus Naderi Zarneh, Jalaluddin TajPour, Saleh Barzegar

*Department of Agronomy, Islamic Azad University, Alborz Branch, Alborz, Iran*

### Abstract

The present article is aimed at study on agroclimatic conditions of cultivation of sunflower throughout selected substations at Kurdistan Province by means of GIS system. In this investigation, meteorological data have been received from synoptic stations based on daily, monthly, and annually trend from Iran Meteorological Organization (IMO) at Kurdistan Province and then homogeneity of data has been explored by (Wald- Wolfowitz) Run Test. Methodology of the research is of statistical descriptive type. Data analysis was carried out by means of Growing Degree Day (GDD) technique and method of Deviation from Optimum Percentage (DOP) plus phenology index as well as thermal potential within environment of statistical software (EXCEL and SPSS). Under agroclimatic conditions, the results of this survey may indicate that the time period among July and September is considered as active months in terms of agriculture based on thermal potential in this region. The In mid-May are the best calendar for cultivation of sunflower month in all the aforesaid substations. Time of harvesting sunflower crop is end August for Bijar, Saghez, and Sanandaj stations while this time is Mid-August for Marivan station. With respect to the phenological method, dates of cultivation until budding, flowering, and the end of flowering stage and maturation start respectively sooner in Marivan station than other stations in this region.

**Keywords:** Agroclimate, Sunflower, Thermal potential, Phenology, Kurdistan


### Introduction

In addition to plant's phenotype, yield of any plant is influenced by climatic conditions (Andrade et al, 2001). Date of cultivation and its density also affected on elements of yield in such a way it was demonstrated that as cultivation is postponed, yield components reduced (Barros, 2004). Appropriate cultivation in various areas may increase photosynthetic efficiency and transfer photosynthetic substances and their storage in seeds and rising yield rather than affecting on rate of germinating and generative growth in plant; at the same time, due to many reasons including high temperature at early period of growth, flowering reduced time, weather coldness, and solar reduced radiations, seed yield is reduced with delayed cultivation (Khalifa, 1984). In addition to determination of appropriate time for cultivation and with selection of suitable density, which is affected by plant's genotype factors and ambient agents

and essential in yield status, the possibility for optimal use of environmental factors and reduction of inter-species competition may be provided (Agele et al, 2007). Sunflower (*Helianthus annuus L.*) as an annual plant from family of *Compositae* or *Asteraceae* and with about 40-50% oil content and high quality and adaptable to various climatic conditions and tolerant to drought may be cultivated under conditions of dry- farming and different irrigation regimes. The potential for production of dry- farmed sunflower under Mediterranean conditions strongly depends on available water supply and the yield ratio of consumption by the given plant Overall, rather than its genotype, the yield of any plant is also influenced by climatic conditions (CERNY et al, 2011). In this regard, (Stone et al, 2001) declared that the rate of access to soil moisture is the paramount factor in determination of yield in farming plants in semi-arid regions. (Pankovich et al, 1999) stated that during budding stage up to the end of flowering phase, shortage of humidity may exert the maxi negative impact on yield of sunflower hybrids. In an experiment, (VARGA, L, 2011) observed that occurrence of stress at graining phase might reduce seed yield but the intensity of this phase does not depend on period of flowering and pollination. (Gomez et al, 1991) noticed that period of germination of sunflower might reduce due to dry stress. Within study on effect of dry stress on some physiological features and elements of sunflower's yield, (LOBELL et al, 2007) reported that water stress has reduced extremely seed yield, biomass, and period of germination period (Bang et al, 1997). The base temperature is about 6°C for growth of sunflower and it grows within mean temperature of 10 to higher than 32°C during day and night. The appropriate growth of sunflower occurs within daily mean temperatures (18-22°C). This plant may initially tolerate well to bud up to -5°C and then it becomes sensitive to coldness and will be frozen under -2°C and lower temperature at 6-foliated stage. Immature seeds may be damaged under icy condition while ripe seeds may be injured less. Sunflower has a developed rhizome, which makes this plant resistant to drought. This plant is resistant to dryness at germination growth stage and at the end of seed maturity, but it is sensitive to drought from stage of observing the calyx up to total coloration (pigmentation) of seeds. Production of sunflower under dry- farming conditions may need to at least 300-350mm precipitation during growth season (GALLIKOVA et al, 2007; Aiken, 2005). Accordingly, any plant may reach to certain period of its growth when it has received certain rate of temperature from its ambience and this is expressed as thermal unit or temperature- day- growth so that this is exclusively a function of genotype (Gilmore and Rogers, 1985). Climatic impact on plant's growth is exerted through acceleration or retardation of growth phases. As a result, growth stages in plants may be predicted with access to the rate of needed temperature for any stage of growth in farming crops and with respect to temperature at studied region.

## Materials and Method

In the current research, the parameters of the maxi and mini daily temperatures have been used relating to statistical period (2002-2011) as well as monthly temperature during statistical career (1991-2011) at Kurdistan Province (**Figure 1**).



**Figure 1:** The studied region

### **Evaluation of thermal potential**

Regarding cultivation of a certain plant, identifying and recognition of thermal potential require using maxi and mini climatic daily data. Temperature serves as one of the main effective factors on growing the plants. With respect to importance of thermal conditions in agriculture and in order to examine thermal characteristics of the region, statistical data for the maxi and mini daily temperatures during 10 years have been explored that they included 36500 cases of daily detection from five selected meteorological substations with their means in Excel software environment from 2001 through 2011.

Afterward, dates of occurrence for the given limits (0 and 10) were characterized and then rates of their occurrence were computed through summation of temperature up to date of occurrence of the same events and presented as some tables. Phenological conditions for any plant species may be measured by identifying and determination of thermal potentials in any region and having these charts and to derive rhythm and trend of their growth. In this case, sum of monthly and annual active temperatures and higher than zero and ten centigrade degree as well as sum of annual active and below- zero temperatures, which are important agriculturally, were calculated and drawn as thermal potential diagrams for any station.

### **Thermal (temperature) Gradient Method**

In order to explore into the studied region in terms of temperature and in relation with rate of deviation from optimum conditions at various heights or optimal time states based on height, it was required adapting thermal gradient technique to determine temperature in height of some points which lacked

measurement substitution. Linear regression method has been utilized to derive these temperatures. By the aid of linear regression, coefficients of temperature variance plus their height have been calculated for months of a year and total year. To compute line equation, the following formula was used:

$(y = ax + b)$  In this formula,  $y$  (independent variable) is the most important variable based on which it is predicted for the expected value (dependent variable)  $x$ . ( $a$ ) denotes a fixed coefficient that is called intercept and ( $b$ ) is slope or thermal (temperature) gradient that represents temperature loss along with height.

The following formulas are employed to calculate  $a$  and  $b$  :

$$a = \frac{\sum(y) \sum(X^2) - \sum(x) \sum(xy)}{N \sum X^2 - (\sum X)^2} \quad (\text{eq.1})$$

$$b = \frac{N \sum XY - (\sum X)(\sum Y)}{N \sum X^2 - (\sum X)^2} \quad (\text{eq.2})$$

To derive the results and calculation of the above formulas, first a table is drawn for correlation among the components for selected substations and the studied time zones formed for each of them so that they will be mentioned as monthly and annual correlation elements for the selected substations.

### **Deviation from Optimum Percentage (DOP) technique**

There are 4 phenological phases in sunflower plant and it has one optimum or optimal temperature per phase where its maxi growth occurs at this optimum temperature. Through identifying and determination of these optimum values for any phenological phase and mean daily temperature derived from detection of mini and maxi daily values, one could characterize spatial optimums within various temporal intervals, particularly months of a year and in fact the points with mini deviation from optimum conditions serve as optimal location. To achieve several spatial optimums in this method, first the optimums or optimal temperatures were determined and then by considering daily statistical mean values, derived difference from the given values about optimum point was computed and at next step, the rate of deviation from optimum conditions were acquired for the above-said locations and their results were identified as tables.

### **Thermal coefficient method or sum of Growing Degree Day (GDD)**

With respect to importance that is attached to temperature cumulative units (degree/day) in identification and topology of appropriate regions for sunflower cultivation and determination of cultivation and harvest dates for this crop based on the given thresholds, Growing Degree Day (GDD) technique has been adapted for this purpose. The above data were processed and analyzed by means of functions in Excel software. In this investigation, the active method (GDD) was used among the common techniques for approximation of thermal units. There are two major techniques for summation of temperature as follows: Sum of effective and active degree day method where sum of active degree day technique has been employed in this study.

#### **a) Sum of active degree day technique**

Phenology or knowledge of phenomena is one of the scientific topics in ecology in which plant's life cycle, which ranged from time of germination to permanent hibernation, is explored.

With respect to climatic variations, especially temperature and soil moisture, dates of start and termination points for each period may differ in several years.

To summate temperature, all diurnal temperature values (without subtracting base temperatures) and during active germination days are added.

The calculation formula is as follows:

$$\frac{T_{Min} + T_{Max}}{2} \quad \text{if} \quad \frac{T_{Min} + T_{Max}}{2} = T_t \quad (\text{eq.3})$$

Where in this formula,  $T_{max}$  and  $T_{min}$  are the maxi and mini daily temperatures and  $T_t$  is biological temperature in this equation. In method of sum of active degree day, which has been also used in this research, sum of daily temperature degrees was used with positive values, but they have been used only for those days in which mean temperatures were higher than biological threshold or biological zero point. All values with quantities greater than  $10^{\circ}\text{C}$  will be calculated while the values with less than  $10^{\circ}\text{C}$  will be excluded from this computation.

### **b) The method determining interval within the stages in phenological studies**

To improve efficiency and properly use from irrigation and implementation of farming operation at any phase of growing the sunflower plant, the needed planning may be executed for growth of crop with determination of the necessary period for both phenological phases based on statistical daily temperature and indentifying interval in the given stage. For this reason, the following formula is used in order to determine the necessary time interval between two phenological phases or (inside stage) based on mini temperature.

$$n = \frac{A}{T - B} \quad (\text{eq.4})$$

**n** denotes the needed time between two phenological phases, **(A)** is thermal coefficient for its completion at the given step, **(B)** as biological threshold of crops, and **(T)** is daily temperature.

## **Findings**

### **The review on thermal potential of the region**

Active temperature degree is a thermal range in which plants are active for growth and germination. With respect to importance of thermal potentials and coefficients in farming activities, sum of annual active and passive temperatures have been studied in above-said height at the level of selected substation. Sum of active temperatures higher than zero ( $0\text{C}^{\circ}$ ) has been computed where many thermal thresholds and biological limits of several plants are situated in this range. Likewise, passive temperature degrees, which represent inertia point for many plant- related activities, were purposed. These achievements may be widely used as framework for future agricultural studies with access to thermal coefficients and the related thresholds. Rhythm and trend of growth in several plant types will be characterized through acquisition and calculation of phenological stages and insertion of the given coefficients in the aforesaid frames. In fact, as the first experience in this field, review on active and passive temperature degrees based on statistical data from maxi and mini daily temperature degrees may be considered as basis for many farming and animal- related activities.

**Table 2:** Sum of monthly and annual active temperature degrees higher than zero and 10C°

Time Station		20 April	21 May	21 June	22 July	22 August	22 September	22 October	21 November	21 December	20 January	19 February	20 March
		Bijar	> 0	Since 21 <sup>st</sup> March (376)	911	1588	2388	3149	3848	4352	Till November 13 <sup>th</sup> (4609)	-	-
Bijar	> 10	-	-	-	388	1113	1891	2410	-	-	-	-	-
Saghez	> 0	-	144	582	1129	1792	2504	3089	3408	-	-	-	-
Saghez	> 10	-	-	-	115	778	1490	2007	-	-	-	-	-
Sanandaj	> 0	-	191	595	1126	1772	2462	2973	3015	-	-	-	-
Sanandaj	> 10	-	-	-	-	646	1336	-	-	-	-	-	-
Marivan	> 0	-	272	732	1333	2055	2836	3426	3787	3842	-	-	-
Marivan	> 10	-	-	-	-	428	1150	1931	2521	-	-	-	-

**Table 3:** Sum of annual active temperature degrees higher than zero and 10C°

Height (m)	Cumulative heat		Σtmean>0	Σtmean>10
	Station			
1883	Bijar		3819	2410
1522	Saghez		3408	2007
1373	Sanandaj		3015	1336
1286	Marivan		3842	2521

To indicate thermal potential status in the studied region, the results, which have been derived based on above tables, were utilized. The results from active and passive temperature degrees based on daily temperature data in 36500 cases of detection show that results of these cumulative temperature or thermal potentials of the region may be deemed as basis for agriculture.

Two thresholds have been considered for this purpose where one is zero- degree threshold in which most of plant and animal- related activities stop and the other threshold belongs to 10C° degree where most of growth activities take place from this threshold and higher level. The results of these calculations show at the level of selected substations that compared to other substations, Marivan station has the maxi cumulative temperature degree higher than zero level (3482) and higher than threshold 10C° (2521C°). Among these stations, Bijar, Saghez, and Sanandaj are respectively considered as the coldest regions of



this province with 4, 4, and 2 monthly active temperature degrees in terms of growth while the maximum active temperature degree occurs in Bijar, Saghez, and Sanandaj substations from early July to mid September.

### Analysis of Deviation from Optimum Percentage (DOP)

Sunflower plant includes four phenological phases, which are important from agroclimatic point of view and reviewed in this investigation. These stages in sunflower are as follows: Cultivation till budding, flowering, end of flowering, and total maturation. Any phase includes an optimum or best temperature in which the plant may grow at maximum level in this optimal temperature. In order to conduct phenological study on sunflower and with respect to the executed investigation, the mid-matured varieties of this crop with the most frequency were considered as base crop. Table (4) shows the rate of deviation from optimum conditions at any phenological stage based on mean daily temperature throughout the selected substations. With respect to derived results from the following table for sunflower plant at flowering stage, compared to other substations, Marivan station has the minimum deviation with higher optimal conditions. Then Bijar, Saghez, and Sanandaj substations have less deviation from this condition.

As a result, compared to other substations, Marivan station has less deviation from optimum status and this means that the aforesaid station possesses optimal conditions for cultivation of sunflower.

**Table 4:** Determining deviation from optimum conditions at phenological phases in sunflower in selected substations

Growth phases Substation	Cultivation to budding		Flowering		End of flowering		Total matured		Sum of deviations
	Optimum	Deviated from conditions	Optimum	Deviated from conditions	Optimum	Deviated from conditions	Optimum	Deviated from conditions	
Bijar	17	-8.64	17.5	-6.24	18	-4.70	20.5	-4.71	-24.28
Saghez	17	-8.19	17.5	-5.92	18	-4.26	20.5	-4.13	-22.50
Sanandaj	17	-10.58	17.5	0.56	18	-5.82	20.5	-5.49	-21.34
Marivan	17	-8.37	17.5	-6.46	18	-4.79	20.5	-4.84	-24.46

### The rate of deviation from optimum conditions based on height

#### Thermal (temperature) gradient

In order to review on rate of deviation from optimum conditions at various heights or spatial optimum conditions based on height, initially coefficients of variance for daily temperatures in respect of height have been calculated for months of a year and total year by means of linear regression technique. To derive the given results and computation of above formulas, firstly correlation elements table was made for the selected substations and in all studied time intervals and a summary of its results has been illustrated as annual correlation elements for the selected substations in Table (5).

**Table 5:** Annual correlation elements of Kurdistan province selected substations during phenological phases (Thermal gradient) for sunflower

Period Coefficients	Cultivation to budding phase	Flowering	End of flowering	Maturation
B	0.002	0.008	0.001	0.001
A	5.30	5.85	3.22	3
R	0.47	0.95	0,32	0.42

Benefitted from regression formula, we calculated thermal gradient table, which denotes status of variable of daily temperature in several heights and months of a year in Excel software environment and by means of the given linear regression regarding the relationship among rate deviation from optimum conditions at any phenological phase and all of its stages and drew its diagram. Due to high  $R^2$ , zoning operation became possible in GIS environment.

### Temporal optimum based on method of active degree days

Active temperature degrees are one of the other agro-climatic methods for determination of optimum times based on the date of latest mini threshold events at any phenological stage (sunflower) that has been used in this investigation. Sum of daily temperatures was used with positive values but only for those days with temperatures, which are higher than average biological level or zero degree of activity. In this study, the basis point for calculation of active thermal coefficients is determined based on two modes: One is based on the mini thresholds of the plant (sunflower) at each of phenological stages and the latter is zero point ( $0C^\circ$ ). Given these plant species extremely depend on temperature so statistical daily temperature has been used as mini and maxi detection data for phenology of plant species (sunflower). Date of completion for each of phenological stages has been determined with identifying accurately each of thresholds in plant's phenological phases (sunflower) and daily temperatures. Date of the mini biological threshold event was considered more than  $10C^\circ$  to activate the plant (sunflower) in each of substations. It requires using 100, 500, 1000, and 1800 thermal units (Btu) higher than zero degree ( $0C^\circ$ ) to achieve date of completion of phenological cultivation stage (sunflower) respectively in each of cultivation stage until budding, flowering phase, at the end of flowering step, and the maturation stage. With respect to Table (6), date of cultivation until budding, flowering, end of flowering, and maturation of sunflower crop start sooner respectively in Marivan, and Saghez substations than other stations.

**Table 6:** Date of completion of phenological stages in sunflower plant

Substation	Height	Date of minithresh old event	Cultivation until budding	Flowering	End of flowering	Total maturation
Bijar	1883	15 <sup>th</sup> May	28 <sup>th</sup> May	22 <sup>th</sup> June	17 <sup>th</sup> July	28 <sup>th</sup> August
Saghez	1522	12 <sup>rd</sup> May	24 <sup>th</sup> May	10 <sup>th</sup> June	8 <sup>th</sup> July	23 <sup>th</sup> August
Sanandaj	1373	17 <sup>th</sup> May	27 <sup>th</sup> May	17 <sup>nd</sup> June	23 <sup>rd</sup> July	26 <sup>th</sup> August
Marivan	1286	5 <sup>st</sup> May	17 <sup>th</sup> May	1 <sup>th</sup> June	3 <sup>th</sup> July	18 <sup>th</sup> August



The date of completion for each of phenological stages is considered as the appropriate method to determine the best cultivation time (sunflower) based on its vital thresholds. The acquired dates are complied with temporal optimums.

### The best cultivation calendar (sunflower)

Determination of cultivation of a crop is considered as one of the objective in agroclimatic studies for which optimum time is obtained for harvesting the crop by surveying climatic conditions. The climatic condition is deemed as the most major factor for harvesting calendar. Among climatic elements, temperature play essential role. With respect to the conducted analyses and based on various agroclimatic methods and phenological studies, the best calendar for crop harvest (sunflower) throughout the studied region that located in different areas are characterized in Table (7).

**Table 7:** Sunflower harvest calendar throughout the selected substations

Station	Date of harvest
Bijar	28 <sup>th</sup> August
Saghez	23 <sup>th</sup> August
Sanandaj	26 <sup>th</sup> August
Marivan	18 <sup>th</sup> August

The computed results and findings are totally complied with comment of experts from farming sector at provincial Agri-jihad department within several areas. In fact according to the conducted interview with farming experts from various towns in the selected substations, they have considered the given climatic calendar as appropriate with respect to regional conditions and potential. Climate plays a determinant role at the beginning period of cultivation and harvest of crop (sunflower).

### Conclusion

Whereas one of the important problems in modern world is production of more foods and nutrients with higher quality and since producing agricultural crops and capabilities of any region depend on its weather and climatic and ambient specifications thus it is crucially important to study on the effective meteorological and environmental on agriculture. Today, Itis deemed as a secured and undeniable platform for accurate development of agriculture. In terms of agroclimatic aspect, sunflower is considered as one of active monthly agricultural crops based on thermal potential method from June to mid October in this region. According to agroclimatic analyses, the best cultivation calendar belongs to the optimal cultivation calendar for sunflower plant in all substations at the end of Mid May. Sunflower is harvested in Saghez, Bijar, and Sanandaj in End August while it is harvested in Marivan station at Mid-August. as a result, in comparison with other substations, Marivan substation has less deviation from optimum conditions; namely, this substation possesses optimal conditions for cultivation of sunflower.

### References

1- Andrade, C. A. D., Constantin, J., Scapim, C. A., Ebraccini, A. D., Angelotti, F., 2001. Effect of weed competition in different spacing upon yield of three common bean (*Phaseolus vulgaris* L.) cultivars. *Ciencia, Eagroecnologica*. 43: 561-568.

- 2- Agele, S. O., Mraiyesa, I. O., Adeniji, I. A., 2007. Effects of variety and row spacing on radiation interception, Partitioning of dry matter and seed set efficiency in late season sunflower (*Helianthus annuus* L.) in a humid zone of Nigeria. African Journal of Agriculture Research. 3: 80-88.
- 3- Barros JFC, Del Carvalho M and Basch G(2004) Response of sunflower ( *Helianthus annus* L) to sowing date and plant density under Mediterranean conditions. Europ Agronomy Journal 21: 347 – 356 .
- 4- Bange, M.P., G.L. Hammer and K.G. Rickert, 1997, Environmental control of potential yield of sunflower in the subtropics, Australian Journal of Agricultural Research, 48(2), 231-40.
- 5- CERNY, I.-PACUTA, V - VEVERKOVA, A., 2011: Uroda a obsah tukov naziek slnecnice rocnej (*Helianthus annuus* L) vplyvom poveternostnych podmienok rocnika a mimokorenej vyzyvy Penatakeepom a Atonikom. In Prosperující olejniný (sborník z konference). Praha: KRV CZU Praha, 118 – 120 s. ISBN 978 - 80 - 213 - 2218 - 9.
- 6- Gomez, D., O. Martinez, M. Arona, and A. Castro. 1991. Generation a selection index for drought tolerance in sunflower. I. Water use and consumption. *Helia*, 14, Nr. 15:65-70.
- 7- Gilmour, E.C. and J.S. Rogers, 1985, Heat unit as a method of measuring maturity in corn. *Agron. J.* 50:611-615.
- 8- GALLIKOVA, M. – KOVACIK, P., 2007: Vplyv sposobu vypoctu davky N hnojiv a terminu ich aplikacie na urodove parametre slnecnice rocnej. *Nitra: SPU*.
- 9- Khalifa, F. M., 1984. Effects of spacing on growth and yield of sunflower under two systems of dry farming in Sudan. *J. Of Agri. Sci.* 103: 213-222.
- 10- LOBELL, D.B. - CAHILL, K.N. – FIELD, C.B., 2007: Historical effects of temperature and precipitation on California crop yields. In *Climatic Change*, 187–203 s.
- 11- Pankovic, D.Z., Sakac. S., Kevrosan, and M. Plesnicar. 1999. Acclimation to long term water deficit in the leaves of two sunflower hybrids: photosynthesis, electron transport and carbon metabolism. *Journal of Experimental Botany.* 50:330, 127 –138
- 12- Stone, L .R.D.E, Goodrum, M.N. Jafar, and A.H.Khan. 2001. Rooting front and water depletion Depths in Grain sorghum and sunflower. *Agron. J.* 1105-1110 .
- 13- VARGA, L., 2011: Listova vyzyva–vyznamny intenzifikacny faktor pri pestovaní poľnohospodárskych plodín.[online].[cit. 2012-25-09]. Dostupne na internete: <http://www.rwaslovakia.sk/storage/file/Listov%C3%A1%20v%C3%BD%C5%BEiva%20RWA%20SLOVAKIA.pdf>.