



Effects of different growing media on growth and flowering of zinnia (*Zinnia elegans*)

Ali Salehi Sardoei^{1*}, Sadegh Shahmoradzadeh Fahraji² and Hadi Ghasemi³

¹ Young Researchers and Elite Club, Islamic Azad University, Jiroft Branch, Iran

² Department of Horticultural Sciences, Islamic Azad University, of Jiroft, Iran.

³ Department of Horticultural Sciences, Shahrood University, of Shahrood, Iran.

ABSTRACT

Zinnia elegans, an annual flowering plant of the genus *Zinnia*, is one of the best known zinnias. Present study was conducted to evaluate the effect of different growing media on the growth and flowering of *Zinnia elegans*. Seven different growth media including coconut compost, silt, soil loam, leaf manure, (leaf manure + silt; 2:1), (coconut compost + soil loam; 1:1) and (leaf manure + soil loam + silt; 1:1:1) were used for growing zinnia. The experiment was laid out in Complete Randomized Design (CRD) giving equal importance to treatments. Number of flowers, length of lateral branches, number of lateral branches, stem diameter, plant height (cm), diameter of flower, vase life (day) and fresh and dry root, shoot and total were determined. The present study confirms the fact that selection of the appropriate medium of growth for potted flowering plants in this case *Zinnia elegans* was very important from aesthetic and marketing point of view. The medium must ensure the production of plants of the required quality on cost effective basis. In the present study, leaf manure produced significantly the maximum vase life and diameter of flower while the maximum vase life and diameter of flower was obtained with mix (coconut compost + soil loam; 1:1).

Keywords: Media, Vase life, Number of flowers, *Zinnia*.

INTRODUCTION

Zinnia is a wonderful summer annual flower which is gaining rapid popularity for its variety of colorful blooms. It belongs to family Compositae. It is native to Mexico and Central America. *Zinnia* flowers exhibit bright, uniform colors, sturdy stems with disease resistant plants and long vase life (Dole, 1999). Increasing awareness of environment-related issues, as well as the need to dispose of and use rising amounts of waste along with the need to reduce the consumption of nonrenewable resources like peat have encouraged the use of composted organic biomass in agriculture. In Iran, *Zinnia* is grown in many types of soils, soil mixtures, or mixtures of organic matter and materials without soil that may include sand, peat, perlite, bark and wood chips, sludge, or composted leaves. The growing media should be porous, uniform in texture, hold sufficient moisture and should be well drained. Commercial mixtures are often used because they are sterilized, ready to use and may even contain some fertilizer (Hochmuth et al., 1996). Different growing media can be used to grow *Zinnia* while the physical and chemical properties

of media, like structure, texture, pH as well as nitrogen, phosphorus and potassium are the dominant factors for the growth and development of plant. Composition and nutritional status of the medium is considered to be helpful for the production of good quality flowering plants with more number of flowers and greater size. No extensive work has been carried out on this flowering plant in Iran, therefore, keeping in view the decorative and aesthetic value of the zinnia, this study was undertaken to determine the effect of different growing media on the growth and flowering of zinnia. Pure soil, coconut compost, leaf manure, silt and soil mix were used to evaluate their effects on the growth and flowering of zinnia to find out the appropriate growing media for this species. Good flower production usually depends upon various factors including the type of growing media used. Growing media is defined as the mean where the roots of cultivated plants grow (Kampf, 2000). Their primordial function is to give support for plant growing (Kampf, 2000; Robert, 2000). Nutrients availability plays a pivotal role in good flower production and thus provision of proper growing media is the pre-requisite for better growth and production of floriculture crops. The plant growing medium must be porous for root aeration and drainage and also capable of water and nutrient retention. Oxygen, of course, is required for all living cells. The coarse-textured media often meet these requirements. Mushroom compost, leaf mold, farmyard manure and other amendments may fulfill these requirements. Brundert and Schmidt (1982) stated that plants with higher water requirements grew more vigorously in leaf mold medium. Fernandez (1984) observed that the plant height and leaf development of foliage plants were best in leaf mold medium. Khan and Khan (1991) reported that the bulb of Dahlia was best developed in the leaf mold. Pasini and Aquila (1989) observed maximum plant height and number of leaves in plants grown in leaf mold medium. Shah *et al.* (2006) got maximum leaves (7.0), with maximum length (20cm), maximum leaf area (84.7cm²) and maximum roots (15) in *Ficus binnendijki* cutting, when leaf mold was used as potting media. A good growing medium would provide sufficient anchorage or support to the plant, serve as reservoir for nutrients and water, allow oxygen diffusion to the roots and permit gaseous exchange between the roots and atmosphere outside the root substrate (Argo, 1998; Abad *et al.*, 2002). This important factor is usually underestimated while working on commercial flower production. Only few studies deal with growing media and their effect in various flowering ornamentals like Liliaceae (Kapoor *et al.*, 2000), gladiolus (Salim *et al.*, 2002), Crocus (Wazir, 2005), Phlox (Naz *et al.*, 2006), Dahlia (Kiran *et al.*, 2007), Freesia (Ali *et al.*, 2011) and tuberose (Ikram *et al.*, 2012). The current study is being proposed to study the effect of different growing media on growth and flowering of Zinnia (*Zinnia elegans*) plant.

MATERIALS AND METHODS

Plant Material and Treatments

Present research was conducted in the Floriculture area, Institute of Horticultural Sciences, University of Agriculture, Jiroft. Seven different growth media including coconut compost, silt, soil loam, leaf manure, (leaf manure + silt; 2:1), (coconut compost + soil loam; 1:1) and (leaf manure + soil loam + silt; 1:1:1) were used for growing zinnia.

Plant-Growth Parameters

Pots of 17.5 cm diameter were used for growing plants. Experiment was carried out in CRD having twenty eight plants in an experimental unit with seven replications and data were collected fortnightly. Thirteen different parameters including number of flowers, length of lateral branches, number of lateral branches, stem diameter, plant height (cm), diameter of flower, vase life (day) and fresh and dry root, shoot and total were used for this study.

Data Analysis

Analysis was performed on data using SPSS 16. Comparisons were made using one-way analysis of variance (ANOVA) and Duncan's multiple range tests. Differences were considered to be significant at $P < 0.05$.

RESULTS AND DISCUSSION

Seven different media used in this study had significant impact on some of the selected parameters of vegetative growth and ornamental value of *Zinnia elegans* the effect was statistically or non significant on Stem diameter, root fresh weight and root and shoot dry weight (Table 1). In general, of leaf manure gave the highest values of growth parameter such as vase life and diameter of flower which were significantly. Growth medium is known to have a large effect on value of potted ornamental plants (Vendrame *et al.*, 2005). The effect of various media with regard to one of the important parameters length of lateral branches and number of lateral branches was significant and the leaf manure mix (leaf manure + silt; 2:1) gave significantly the highest values as compared to all other media which differed significantly with one another (Table 1).

As far as the visual qualitative assessment of flowers was concerned, the differences due to media were or non significant (Table 2) and, therefore, the role of the media tested seems to be unimportant in this regard. The effect of various media with regard to one of the important parameter number of flowers per plant was significant and the leaf manure mix (coconut compost + soil loam; 1:1) gave significantly the highest values as compared to all other media which differed significantly with one another (Table 1). It is worth mentioning that the leaf manure had produced most vigorous plants in terms of vase life, diameter of flower and root and total fresh and dry weight compared with other media. Similar results were reported by Riaz *et al.* (2008) who also reported.

Table 1- Effect of Various Growth Media on Growth and Flowering Characteristics of *Zinnia elegance*

Media	Stem diameter (cm)	Plant height (cm)	Length of lateral branches (cm)	Number of flowers	Number of lateral branches	Vase life (day)	Diameter of flower (mm)
soil loam	0.22a	44.50cd	19.03c	3.3abc	2.25b	27.25ab	63.45bc
leaf manure	0.31a	49.37bcd	24.35bc	3.5ab	3ab	32.75a	72.27a
coconut compost	0.26a	62.25a	31.23ab	3.25abc	3.25a	26.50ab	66.58ab
leaf manure + silt	0.23a	54.37abc	36.37a	3.5a	3.5a	24.25ab	65.85ab
leaf manure + soil loam + silt	0.23a	58.12ab	28.62abc	2.75bc	2.25b	32.25a	69.71ab
coconut compost + soil loam	0.26a	42.87d	22.47bc	3.75a	2.75ab	20.50b	67.52abc
silt	0.26a	50.87bcd	24.11bc	2.5c	2.75ab	23.33ab	61.11c

Means followed by same letter are or non significantly different at $P < 0.05$ probability using Duncan's test.

These results support the findings of Awang & Ismail (1997) and Riaz *et al.*, (2008) who found that *Zinnia elegans* produced more flowers where grown in a medium containing coconut coir although, in this case, coconut coir alone failed to produce these effects. However, the coconut compost alone produced significantly the largest average plant height (62.25 cm) compared with rest of the media which were statistically at par with one another, although the leaf manure mix was next in the order to the coconut compost in terms of average plant height (Table 1). Similar results were reported by Rahbarian and Salehi Sardoei (2013) who also reported. The results showed that the availability of proper nutrients to the plants may alter the plant height as Younas *et al.* (2008) also obtained maximum plant height of *Dahlia coccinia* on leaf manure. Similar results have been reported by Riaz *et al.* (2008) who reported vigorous plant height in *Zinnia elegans* cv. Blue point, when Leaf manure mixture was used as a growing media. The effect of various media with regard to one of the important parameters shoot fresh and dry weight was significant and the leaf manure mix (leaf manure + silt; 2:1) gave significantly the highest values as compared to all other media which differed significantly with one another (Table 2).

Table 2- Effect of Various Growth Media on Plant Fresh and Dry Weight of *Zinnia elegans*

Media	Fresh weight (g)			Dry weight (g)		
	Root	Shoot	Total	Root	Shoot	Total
soil loam	1.78a	13.72ab	15.50c	0.99a	6.78a	7.78bc
leaf manure	3.82a	31.05a	35.02a	2.12a	15.89a	18.01a
coconut compost	2.67a	21.52a	24.19b	1a	11.69a	12.69ab
leaf manure + silt	3.07a	31.41a	34.48a	1.33a	16.59a	17.93a
leaf manure + soil loam + silt	3a	25.42ab	28.42ab	1.42a	11.32a	12.74ab
coconut compost + soil loam	2.25a	23.15ab	25.41b	1.16a	12.05a	13.22a
silt	1.01a	11.15b	12.56c	0.55a	4.86a	5.41c

Means followed by same letter are or non significantly different at $P < 0.05$ probability using Duncan's test.

CONCLUSIONS

The present study confirms the fact that selection of the appropriate medium of growth for potted flowering plants in this case *Zinnia elegans* was very important from aesthetic and marketing point of view. The medium must ensure the production of plants of the required quality on cost effective basis. In the present study, leaf manure produced significantly the maximum vase life and diameter of flower while the maximum vase life and diameter of flower was obtained with mix (coconut compost + soil loam; 1:1) (Rahbarian and Salehi Sardoei, 2014a, b; Salehi Sardoei and Rahbarian, 2014).

REFERENCES

- Abad, M., Noguera, P., Puchades, R., Maquieira, A and Noguera, V (2002). Physico- chemical and chemical properties of some coconut dusts for use as a peat substitute for containerized ornamental plants. *Biores. Technol.* 82: 241-245.
- Ali, T., Khattak, A.M. Amin, N.U and Khan, M.A. (2011). Response of freesia cultivars to different growing media under Peshawar conditions. *Sarhad J. Agric.* 27: 43-49.
- Riaz, A., Arshad, M., Younis, A., Raza, A and Hameed, M (2008). Effects of different growing media on growth and flowering of *Zinnia elegans* cv. Blue point. *Pak. J. Bot.* 40(4): 1579-1585.
- Awang, Y and Ismail, M (1997). The growth and flowering of some annual ornamentals on coconut dust. *Acta Hort.* 450: 31-38.
- Argo, W.R (1998). Root medium chemical properties. *Horticultural Technology.* 8: 846-894.
- Brundert, W and Schmidt, K (1982). Organic composts in hydroculture and watering method. *Dentscher Gertenban* 36: 904-906.
- Dole, H.C (1999). Zinnias: Colorful, Butterfly-Approved. *Butterfly Gardeners Quarterly.* BGQ, PO Box 30931, Seattle, WA 98103.
- Fernandez Maldonado, J.F (1984). Trials on different substrates for ornamental plants. *Bolitin informatvo, estacion de investigation sobre cultivos. Horticolas intensivos* 5: 89-107.
- Hochmuth, R.C., Davis, L.L and Dinkins, D (1996). Greenhouse cut flower zinnia production Florida Cooperative Extension Service. 3(48): 98-102.
- Ikram, S., U. Habib and N. Khalid. 2012. Effect of different potting media combinations on the growth and vase life of tuberose. *Pak. J. Agri. Sci.* 49: 121-125.
- Waseem, K., Hameed, A., Saleem Jilani, M., Kiran, M., Rasheed, M.U., farullah, G., Javeria, S and Ali Jilani, T (2013). Effect of Different growing Media on the Growth and Flowering of Stock (*Matthiola incana*) under the Agro-Climati Conditation of Dera Ismail Khan. *Pak. J. Agri. Sci.* 50(3): 523-527.
- Kampf, A.N (2000). The substrate Commercial production of ornamental plants. *Guaíba: Agriculture.* pp. 254.
- Kapoor, M., Grewal, H.S and Aror, J.S (2000). Effect of media on propagation of Liliun. *J. Ornamental Hort. New Series.* 3: 58-59.
- Khan, M.A and Khan, A.B (1991). "Dahlia". *Zarat Nama.* 29: 9-11.
- Kiran, M., Baloch, J.U.D., Waseem, K., Jilani, M.S and Khan, M.Q (2007). Effect of different growing media on the growth and development of Dahlia (*Dahlia pinnata*) under the agro- climatic condition of Dera Ismail Khan. *Pak. J. Biol. Sci.* 10: 4140-4143.

- Naz, I., Baloch, J.U.D and Waseem, K (2006). Effect of different growing media on the growth and development of Phlox (*Phlox drummondii*). M.Sc (Hons) Thesis. Department of Horticulture, Faculty of Agriculture, Gomal University, D.I.Khan.
- Pasini, C. and Aquila, F.D (1989). Trials on the cultivation of Diffenbachia cv. Alix in substrate based on leaf mold and other components. Annali dell institute sperimentale per la floricultura 18:7-9.
- Rahbarian, P and Salehi Sardoei, A (2014a). Effect of different Media on growth, Sucker and Chlorophyll of *Padanus spp* in Under System Mist. International Journal of Farming and Allied Sciences. 3(3): 285-288.
- Rahbarian, P and Salehi Sardoei, A (2014b). Effect Waste of palm trees and sand and perlite mixed with some growth indices *Ficus Benjamina*. International journal of Advanced Biological and Biomedical Research. 2(3): 573-578.
- Rahbarian, P and Salehi Sardoei, A (2013). Effect of Different Substrates on *Rosmarinus officinalis* L. Production in Arrigation Mist. Journal of plant ecophysiology. In press.
- Robert, R. (2000). Horticultural substrates: possibilities and limits of its composition and use, examples of research, industry and consumption: Substrates for plants the basis of plant production in containers. Port Alegre: Genesis. pp. 209-215.
- Salehi Sardoei, A and rahbarian, P (2014). Effect of Different Media on Growth Indexes of Ornamental Plants Under System Mist. International European Journal of Experimental Biology. In press.
- Salim, K., Hameed, A.K., Aqib, I and Rehman, J.M (2002). Effect of different media on growth and quality of gladiolus (*Gladiolus hortulanus* cv. Jacksonville Gold). Asian J. Pl. Sci. 1:670-671.
- Shah, M., Khattak, A.M and Amin, N.U (2006). Effect of different growing media on the rooting of *ficus binnendijkii* amstel queen cuttings. J. Agric. Biol. Sci. 1:15-17.
- Wazir, K.U. 2005. Response of crocus (*Crocus spp.*) cultivars to different growing media. M.Sc. Thesis, Agric. Univ. Peshawar, Pakistan.
- Younas, A., Ahmad, M., Riaz, A and Khan, M.A (2008). Effect of different potting media on growth and flowering of *Dahlia coccinia* cv. mignon. Acta Hort. 804: 191-196.